



IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY



VOLUME : 5 ISSUE : 2 Print / Issue Publication Date: 12-Aug-2020



ISSN : 2455-2143



DOI : 10.33564/IJEAST.2020.v05i02.003

Indexed In



WWW.IJEAST.COM

editor@ijeast.com



EVALUATION OF SAPONIFICATION VALUE, IODINE VALUE, PEROXIDE VALUE AND FREE FATTY ACID LEVEL OF ESSENTIAL OIL OF CAYENNE PEPPER (*Capsicum annum*)

Aguebor-Ogie Nogiomwan Bobby

Department of Medical Biochemistry, SBMS, CMS
University of Benin, Benin City, Edo State, Nigeria

Nwosu Favour Ihuoma

Department of Medical Biochemistry, SBMS, CMS
University of Benin, Benin City, Edo State, Nigeria

Edokhume Peter

Department of Medical Biochemistry, SBMS, CMS
University of Benin, Benin City, Edo State, Nigeria

Abstract— Cayenne peppers (*Capsicum annum*) have been shown to be the world's most cultivated and consumed spice, especially in the temperate and semi-temperate regions respectively. It has also been known for its antioxidant, anti-mutagenic and hypocholesterolemic properties as well as it containing essential oil. Here, the total flavonoid and phenol contents and essential oil of the pepper were evaluated for its physiochemical properties which included iodine value, saponification value and free fatty acid value and peroxide value. The essential oil was extracted and the physiochemical properties were analyzed. The essential oil was shown to contain free fatty acid ($0.30 \pm 0.04\%$), iodine value ($23.88 \pm 0.23\%$), peroxide value ($1.23 \pm 0.02\%$) and Saponification value ($141.10 \pm 0.35\%$). The result indicated that, continued usage of the pepper as a major condiment in preparation of some foods (soups and/or sauces), allows for its medicinal importance to be exploited maximally in relation to health.

Keywords— Flavonoid, Phenol, Cayene pepper, essential oil

I. INTRODUCTION

Capsicum annum (*C. annum*) is a plant's fruit in the genus, *Capsicum*, of the family of nightshade, *Solanaceae*. It has a characteristic pungent odour/scent. It can be cultivated as a value-added processed product, vegetable and spice [1]. It is a common constituent of cooked and/raw food(s). It adds flavour, colour, vitamins (e.g. A and C) and a characteristic aroma. It has become indispensable to so many Nigerian chefs, for home use and food industries worldwide [2] as a result of the presence of these nutrients including flavonoids and phenols that are actively involved in promoting health [3]. In other study, it was reported to have antioxidant, anticancer, analgesic and antipyretic properties [4, 5]. *C. annum* has been shown to

be a very rich source of vitamins A, C, B₆, folic acid and beta-carotene, which is known to provide excellent nutrition for humans [4]. It is being used by security agencies to produce tear gas which is used mostly to control crowds in order to minimize violence [6]. Agronomically, in Egypt, genotypes of various pepper species were found to exhibit differential responses to the Egyptian broomrape which is a chlorophyll-lacking root-parasite and this has led to the use of *C. annum* as a catch/trap crop to minimize infestation of the parasite [7]. Even though the pepper is very popular in all the agro-ecological zones of Nigeria and features amongst the most widely used peppers in typical Nigerian dishes, very little has been achieved in the knowledge of its important chemical constituents, nutritional constituents and medicinal benefits to health. Very limited literatures on the phytochemical properties of locally grown cayenne pepper in Nigeria are available. Therefore, this study aimed to examine the physiochemical properties of Cayenne pepper locally grown in Nigeria.

II. MATERIALS AND METHODS

A. Preparation of plant extract

Fresh fruits of *C. annum* were purchased from a local market (New Benin market) in Benin City, Nigeria and identified at the Plant Biology and Biotechnology Department, Faculty of Life Sciences, University of Benin, Nigeria. They were washed, dried and diced into bits to increase the surface area after which they were oven-dried at 40°C until a dry constant weight was obtained and then pulverized. This (250grams of sample) was then subjected to Soxhlet extraction with n-hexane for 6 hours. Thereafter, the extract containing the n-hexane and the oil were evaporated to separate the n-hexane from the oil.



B. DETERMINATION OF FREE FATTY ACID [8]

The free fatty acids is the estimated by titrating it against potassium hydroxide (KOH) using phenolphthalein as indicator. The acid value is mg KOH required to neutralize the free fatty acids present in one gram of sample. It is expressed as oleic acid (octadec-9-enoic acid) equivalent. Dissolve 10g of oil in 50ml of the neutral solvent (25ml ether plus 25ml 95% ethanol and 1ml phenolphthalein solution neutralized with 0.1M KOH) in 250ml conical flask. Then added 3-4 drops of phenolphthalein indicator and mixed by shaking. Titrated the content against 0.1M KOH until a pink color which persisted for fifteen seconds was obtained.

C. DETERMINATION OF IODINE VALUE OF OIL [8]

A 0.001kg of oil sample was weighed into a 500mL volumetric flask. 15mL of carbon tetrachloride was added to the sample and swirled to ensure that the sample is completely dissolved. 25mL of Wijs solution was then dispensed into the flask containing the sample using a pipette. The flask was stoppered and swirled to ensure complete mixing. The sample was then placed in the dark for 30 minutes at room temperature. The flask was removed from storage and 20mL of 10% potassium iodide (KI) solution added, followed by 150mL of distilled water. The mixture was titrated with 0.1N thiosulphate ($\text{Na}_2\text{S}_2\text{O}_3$) solution, adding gradually and with constant and vigorous shaking until the yellow colour had almost disappeared. 1.5mL of starch indicator solution was added and the titration was continued until the blue colour disappeared. A blank determination was conducted simultaneously.

D. SAPONIFICATION VALUE [8]

A 0.002kg of the oil sample was weighed into a volumetric flask. Then 25mL of 1.0N alcoholic KOH was pipetted and allowed to drain for about 1 minute into the mixture. A condenser was connected to the flask and the mixture sample was allowed to boil gently but steadily for 45 minutes for complete saponification. The flask and the condenser were then cooled but not sufficiently to form a gel, the inside of the condenser was washed down with about 1ml of distilled water. The condenser was disconnected and 1ml of phenolphthalein indicator added. The solution was titrated with 0.5N hydrochloric acid (HCl) until the pink colour just disappeared. A blank determination was conducted simultaneously with the sample.

E. PEROXIDE VALUE [8]

Rancidity is brought about by the action of air (oxidative) or by microorganisms in oil. In oxidative rancidity oxygen is taken by the oil or fat with the formation of peroxides. Peroxide value is a measure of the peroxides contained in the oil. The peroxides present are determined by titration against

thiosulphate in the presence of KI using starch as an indicator. Weighed 1gram of oil into a clean dry boiling tube and added 1gram of powdered KI and 20ml of solvent mix. Transferred the tube into boiling water so that the liquid boils vigorously for not more than 30 seconds. Then transferred the contents quickly to a conical flask containing 20ml 5% KI solution. Washed the tube twice with 25ml of water each time and collect into the conical flask. Titrated with 0.002M $\text{Na}_2\text{S}_2\text{O}_3$ solution to a pale yellow. Further added 0.5ml of starch, shook vigorously and titrated once more, carefully, till the blue color disappeared.

F. Statistical analysis

Statistical analysis for comparing the data was performed by using IBM SPSS Statistical software, version 20 (IBM Corp, Pakistan). Results are showed as Mean $\pm$ SEM by using the analysis of variance.

III. RESULT

The physicochemical properties of oil were determined Table below shows the results from the characterization of the *capsicum annuum* essential oil. It involves the value of the free fatty acid, iodine value, peroxide value and Saponification value.

Table - 1 Saponification Value, Iodine Value, Peroxide Value and Free Fatty Acid Level of *Capsicum annuum*

Parameter	Free fatty acid (mEq/L)	Iodine Value (mgKOH/g)	Peroxide (mgEq O_2 /g)	Saponification (mgKOH/g)
Estimated value	0.30 $\pm$ 0.05	23.88 $\pm$ 0.23	1.23 $\pm$ 0.02	141.10 $\pm$ 0.35

*Values are expressed as Mean $\pm$ SEM. All analyses were done in triplicate.

IV. DISCUSSION

Essential oils or "essences" owe their name to their flammability [9]. Essential oils are soluble in alcohol, ether, and fixed oils, but insoluble in water. These volatile oils are generally liquid and colorless at room temperature. They have a characteristic odor, are usually liquid at room temperature and have a density less than unity. Iodine values are often used to determine the amount of unsaturation in fatty acids. In general, higher is the iodine value, higher is the unsaturation and hence higher is the chances of oxidation upon exposure to atmosphere (i.e., rancidity tendency increases). So, it reduces the shelf life of oil. Lower iodine value means they are non-drying and hence their self life is more. Thus, the iodine value



obtained in this study classify the *Capsicum annum* oils as non drying oils which is in corroboration with earlier work by [10]. [11] reported that high saponification values of fats and oils are due to the predominantly high proportion of shorter carbon chain lengths of the fatty acids. Thus, the estimated saponification value in this study which was high, though not up-to the required range accepted in soap industry, revealed that it contains reasonable level of short chain fatty acids and as well, not too suitable for soap industry. Detection of peroxide gives the initial evidence of rancidity in unsaturated fats and oils among other methods that are available, but peroxide value is the most widely used [12]. It gives a measure of the extent to which an oil sample has undergone primary oxidation [14]. The peroxide value from this study show that the oil could be of pharmaceutical value. Free fatty acids are considered as defect in oils/ fats because they are degraded or become rancid. Presence of free fatty acid causes rancidity in oils. Acid value determination is often used as a general indication of the condition and edibility of the oil. This is because an increase in acid value is accompanied by development of objectionable flavours and odours [15]. Acid value is used as an indicator for edibility of oil and suitability for use in the paint industry [15].

V. CONCLUSION

The preliminary physiochemical studies carried out on *C. annum* essential oil revealed the presence of some secondary metabolites and properties. The essential oil was also found to possess some degree of unsaturation. Some of these properties of cayenne pepper are of great health benefits such as reducing pain, reducing risk of cancer, aids in digestion. Besides the strict medicinal application, cayenne pepper essential oil can be used in other industries such as food industry, cosmetic industry and chemical industry.

VI. REFERENCE

- [1] Kumar, S., and Rai, M. (2005) Chile in India. Chile Pepper Institute Newsletter (XXII). (pp 1-3).
- [2] Dagnoko, S., Yaro-Diarisso, N., Sanogo, P.N, Adetula, O., Dolo-Nantoume, A., Gamby, T.K., Traore-Thera, A., Katile, S., and Diallo-Ba, D. (2013) Overview of pepper (*Capsicum* spp.) breeding in West Africa. African Journal of Agricultural Research. Vol 8, No 13 (pp1108-1114).
- [3] Ogunlade, I., Alebiosu, A.A, and Osasona, A.I. (2012) Proximate, mineral composition, antioxidant activity and total phenolic content of some pepper varieties. International Journal of Biological and Chemical Sciences. Vol 6 No 8 (pp2221-2227).
- [4] Nadeem, M., Muhammad, F., Anjum, M., Khan, A.M.R., Saed, M, Riaz, A. (2011) Antioxidant Potential of Bell Pepper (*Capsicum annum* L.). Pakistan Journal of Food Science. Vol 21 No 14 (pp 45-51).
- [5] Pickersgill, B. (1997) Genetic resources and breeding of *Capsicum* spp. Euphytica. Vol 96 (pp 129-133).
- [6] Smith, C.G. and Stopford, W. (1999) Health hazards of pepper spray. N. C. Med. J. vol 60 No 5 (pp 268-274)
- [7] Hershenhorn, J., Goldwasser, Y., Plakhine, D., Herzlinger, G., Golan, S., Russo, R. and Kleifeld, Y. (1996) Role of pepper (*Capsicum annum* L) as a trap and catch crop for control of *Orobancha egyptiaca* and cernual. Weed Science. Vol 44 (pp 948-951).
- [8] A.O.C.S. (1993) Official Methods and Recommended Practices of the American Oil Chemists Society, 4th Ed. Published by the American Oil Chemists Society, 1608, Broadmoor Drive, Champaign, Illionis. (pp 61826-3489)
- [9] Kyari, M..Z. (2008) Extraction and Characterization of Seed Oils. International Agrophysics. Vol 22 (pp 139-142).
- [10] Nielson, S.S. (1994) Introduction to the chemical analysis of foods. Chapman and Hall, New York. (pp 93-207).
- [11] Kirk, R. and Sawyer, R. (1999) Pearson's composition and analysis of foods. 9th edition, Addison Wesley longman ltd. England. (pp. 9-29, 608-640).
- [12] Dobarganes, M. and Velasco, J. (2002) Analysis of lipid hydroperoxides. European Journal of Lipid Science and Technology. Vol 104 (pp 420-428).
- [14] Gbalade, O. and Soremekun, R.O. (1997) Evaluation of quality of Nigerian chillies for pharmaceutical formulations. Journal of Pharmaceutical and Biomedical Analysis. Vol 15 (pp 545-548).
- [15] Nwokem, C.O., Agbaji, E.B., Kagbu, J.A. and Ekanem EJ. (2010) Determination of capsaicin content and Pungency Level of Five Different Peppers Grown in Nigeria. New York journal. Vol 3 No 9 (pp 17-21).

IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

ABOUT IJEAST

International Journal of Engineering Applied Science and Technology (IJEAST) is a peer-reviewed, open access journal that publishes high-quality research papers in the field of Engineering, Applied Science and Technology.

IJEAST aims to provide a platform for researchers, academicians, and professionals to share their innovative ideas, research findings, and practical experiences with the global scientific community.

FOCUS AREAS

- Engineering
- Applied Science
- Technology
- Innovation & Development
- Interdisciplinary Studies



PEER REVIEWED

All submissions are rigorously peer reviewed to ensure quality.



OPEN ACCESS

Free and unrestricted access to research for all.



GLOBAL REACH

Connecting researchers and professionals worldwide.



TIMELY PUBLICATION

We ensure a swift and efficient publication process.



For more information, visit our website
www.ijeast.com



INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

✉ editor@ijeast.com

🌐 www.ijeast.com

📍 India



2455-2143