

IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

VOLUME : 7 ISSUE : 03 Print / Issue Publication Date: 29-Aug-2022

ISSN : 2455-2143

DOI : 10.33564/IJEAST.2022.v07i03.018

Indexed In

Google SCHOLAR OPEN ACCESS doi Crossref

WWW.IJEAST.COM

editor@ijeast.com

International Journal of Engineering Applied Sciences and Technology, 2022
Vol. 7, Issue 3, ISSN No. 2455-2143, Pages 120-121
Published Online July 2022 in IJEAST (http://www.ijeast.com)

ON NON-HOMOGENEOUS TERNARY CUBIC DIOPHANTINE EQUATION

$$w^2 - z^2 + 2wx - 2zx = x^3$$

S.Vidhyalakshmi
Assistant Professor,
Department of Mathematics
Shrimati Indira Gandhi College,
Affiliated to Bharathidasan University, Trichy-620 002,
Tamil Nadu, India.

M.A.GOPALAN
Professor,
Department of Mathematics,
Shrimati Indira Gandhi College,
Affiliated to Bharathidasan University, Trichy-620 002,
Tamil Nadu, India.

J. Shanthi, T. Mahalakshmi
Assistant Professor,
Department of Mathematics,
Shrimati Indira Gandhi College,
Affiliated to Bharathidasan University, Trichy-620 002,
Tamil Nadu, India.

***Abstract*— The non-homogeneous ternary cubic Diophantine equation $w^2 - z^2 + 2wx - 2zx = x^3$ is analyzed for its patterns of non-zero distinct integral solutions. A few relations between the solutions and special number patterns are presented.**

***Keywords*— Ternary cubic Non- Homogeneous cubic, Integral solutions.**

Notations:

$$t_{m,n} = n(1 + \frac{(n-1)(m-2)}{2})$$

$$P_n^r = \frac{n(n+1)(n(r-2)-(r-5))}{6}$$

$$CP_k^4 = \frac{4k^3 + 2k}{6}, \quad CP_k^8 = \frac{8k^3 - 2k}{6}$$

I. INTRODUCTION

The Diophantine equation offers an unlimited field for research due to their variety [1-3]. In particular, one may refer [4-15] for cubic equations with three unknowns. This communication concerns with yet another interesting equation $w^2 - z^2 + 2wx - 2zx = x^3$ representing non-homogeneous cubic with three unknowns for determining its infinitely many non-zero integral points. A few relations between the solutions and special number patterns are presented.

II. METHOD OF ANALYSIS

The given non-homogeneous ternary cubic diophantine equation is

$$w^2 - z^2 + 2wx - 2zx = x^3 \quad (1)$$

On completing the squares,(1) is written as

$$P^2 - Q^2 = x^3 \quad (2)$$

Where

$$P = w + x, Q = z + x \quad (3)$$

Write (2) as the system of double equations as below:

$$P + Q = x^3,$$

$$P - Q = 1$$

Solving the above system of equations, one obtains

$$x = 2k + 1 \quad (4)$$

and

$$P = 4k^3 + 6k^2 + 3k + 1, Q = 4k^3 + 6k^2 + 3k \quad (5)$$

International Journal of Engineering Applied Sciences and Technology, 2022
Vol. 7, Issue 3, ISSN No. 2455-2143, Pages 120-121
Published Online July 2022 in IJEAST (http://www.ijeast.com)

In view of (3),we have

$w = 4k^3 + 6k^2 + k, z = 4k^3 + 6k^2 + k - 1$ (6)

Thus,(4) and (6) represent the integer solutions to (1).

Relations between the solutions and special number patterns:

I. Each of the following expressions is a square multiple of 2

(i). $w - x * t_{3,2k}$

(ii). $z + 1 - x * t_{3,2k}$

(iii). $w + t_{3,2k} - 3 * P_{2k}^3$

(II). $w = -12t_{3,k} + 24 * P_k^3 - k$

(III). $w = -t_{14,k} + 24 * P_k^3 - 12k$

(IV). $w - x = t_{6,k} + 8 * P_k^5 - 1$

(V).Each of the following is a square multiple of 6

(i). $w - 6CP_k^4 + k$

(ii). $w + z + x - 6CP_k^8 + 12t_{3,k}$

For simplicity and brevity, the other choices of solutions to (1) are exhibited below:

Choice 1:

$x = k, z = t_{3,k-1} - k, w = t_{3,k} - k$

Choice II:

$x = 2na, z = 2n^3a^2 - (2n+1)a, w = 2n^3a^2 - (2n-1)a$

Choice III

$x = 2na, z = a^2 - 2(n^3 + n)a, w = a^2 + 2(n^3 - n)a$

Choice 1V:

$x = 2na, z = 2n^3a^3 - (2na+1), w = 2n^3a^3 - (2na-1)$

Choice V:

$x = 2na, z = a^3 - 2n(n^2 + a), w = a^3 + 2n(n^2 - a)$

III.CONCLUSION

In this paper, we have made an attempt to obtain all integer solutions to (1). To conclude, one may search for integer solutions to other choices of ternary cubic diophantine equations.

IV. REFERENCE

[1]. Dickson,L.E;(1952) History of Theory of Numbers, vol 2, Chelsea publishing company, New York.

[2]. Mordell, L.J;(1969)Diophantine Equations, Academic press, London..

[3]. Carmichael, R.D;(1959),The theory of numbers and Diophantine analysis, New York, Dover.

[4]. Gopalan,M..A;,Srividhya,G.(2011),Integralsolutionsoft ernarycubicdiophantineequation$x^3+y^3=z^2$ Acta Ciencia Indica, Vol XXXVII, No.4,(Pp 805-808).

[5]. Gopalan,M.A;Vidhyalakshmi,S;Thiruniraiselvi,N,(2013), On Non-Homogeneous Cubic Equation With Three Unknowns $x^2 - y^2 + z^2 = 2kxyz$,BOMSR,Vol.1(1), (Pp 13-15).

[6]. Gopalan. M. A; Vidhyalakshmi. S; Lakshmi. K, (2013), Latic points on the non-homogeneous cubic equation $x^3 + y^3 + z^3 + (x + y + z) = 0$, Impact.J.Sci. Tech, Vol.7, No. 1, (Pp 21-25).

[7]. Gopalan. M. A; Vidhyalakshmi. S; Lakshmi. K, (2013), Latic points on the non-homogeneous cubic equation $x^3 + y^3 + z^3 - (x + y + z) = 0$, Impact.J.Sci. Tech, Vol.7, No. 1, (Pp 51-55).

[8]. Vidhyalakshmi,S ;Usharani,T.R;Gopalan,M.A,(2014), Integral Solutions Of The Ternary Cubic Equation $5(x^2 + y^2) - 9xy + x + y + 1 = 35z^3$,IJRET,Vol:0 3,Issue:11,(Pp 449-452).

[9]. Gopalan,M.A; Thiruniraiselvi,N;Sridevi,R,(2015), On The Ternary Cubic Equation $5(x^2 + y^2) - 8xy = 74(k^2 + s^2)z^3$,IJMRME, Vol.1(1),(Pp 317-319).

[10]. Gopalan. M. A; Vidhyalakshmi. S; Shanthi. J; Maheswari.J, (2015), On ternary cubic Diophantine equation $3(x^2 + y^2) - 5xy + x + y + 1 = 12z^3$, IJAR, Volume 1, Issue 8, (Pp 209-212).

[11]. Janaki,G and Saranya ,P;(2016), On theTernary Cubic Diophantine Equation $5(x^2 + y^2) - 6xy + 4(x + y) + 4 = 40z^3$, International Journal of Science and Research- online, Vol 5, Issue3, (Pp 227-229).

[12]. Gopalan,M.A, Sharadha Kumar,(2018), "On the non-homogeneous ternary cubic equation $3(x^2 + y^2) - 5xy + x + y + 1 = 111\ z^3$ International Journal of Engineering and Techniques, 4(5),(Pp105-107).

[13]. Sharadha Kumar, Gopalan,M.A; (2019),"On The Cubic Equation $x^3 + y^3 + 6(x + y)z^2 = 4w^3$",JETIR, 6(1), (Pp 658-660).

[14]. Vijayasankar,A; Dhanalakshmi,G; Sharadha Kumar, Gopalan,M.A;(2020), On The Integral Solutions To The Cubic Equation With Four Unknowns $x^3 + y^3 + (x + y)(x - y)^2 = 16zw^2$ International Journal For Innovative Research In Multidisciplinary Field, 6(5), (Pp 337-345).

[15]. Vijayasankar,A; Sharadha Kumar , Gopalan,M.A,(2021), "On Non-Homogeneous Ternary Cubic Equation $x^3 + y^3 + x + y = 2z(2z^2 - \alpha^2 + 1)$ International Journal of Research Publication and Review2(8),(Pp 592-598).

IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE AND TECHNOLOGY

AUTHORS

- S.Vidhyalakshmi
- M.A.GOPALAN
- J. Shanthi
- T. Mahalakshmi

ABOUT IJEAST

International journal of Engineering Applied Science and Technology (IJEAST) is a peer-reviewed, open access journal that publishes high - quality research paper in the field of Engineering, Applied Science and Technology.

IJEAST aims to provide a platform for researchers, academicians, and professionals to share their innovative ideas, research findings, and practical experiences with the global scientific community.

JOURNAL METRICS

H-INDEX	i10-INDEX	IJEAST CITATIONS	PEER-REVIEWED & OPEN ACCESS
33	154	9,647 (IJEAST CITATIONS)	PUBLISHED MONTHLY

PEER REVIEWED

All submission are rigorously peer reviewed to ensure quality.

AUTHORS ARE OUR PRIORITY

We value our authors and recognize their contribution to the advancement of research and knowledge.

OPEN ACCESS

Free and unrestricted access to research for all.

GLOBAL REACH

Connecting researchers and professionals worldwide.

TIMELY PUBLICATION

We ensure a swift and efficient publication process.

For more information, visit our website
www.ijeast.com

editor@ijeast.com
www.ijeast.com
India

2455-2143