



IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY



VOLUME : 3 ISSUE : 02 Print / Issue Publication Date: 07-Sep-2018



ISSN : 2455-2143



Indexed In



WWW.IJEAST.COM

editor@ijeast.com



QUADRUPLE OF SELF MAPS IN METRIC SPACE AND COMMON FIXED POINT THEOREMS

Rashmi Rani

Department of Mathematics

A.S. College for Women, Khanna, Punjab, India

Bindia Tarika

Department of IT

Punjab Technical University, Jalandher

Abstract— Using property (E.A.) and its variants, we prove some common fixed point theorems for quadruple of weakly compatible self maps in metric space in this paper. Our results extend and unify various known results in literature. We also give application of proved result for four finite families of maps.

Keywords— Metric space, property (E.A), common property (E.A), $JCLR_{ST}$ property, common fixed point, coincidence point.

I. INTRODUCTION

In 2011, Rao and Pant [1] utilized the concept of finite metric spaces and proved some common fixed point theorems for asymptotically regular maps. Recently, Mishra et. al. [2] proved some common fixed point theorems using property (E.A.) (which was introduced by Aamri and Moutawakil [3]) in metric spaces.

Using property (E.A.) and its variants, we prove some common fixed point theorems for quadruple of weakly compatible self maps in metric space in this paper. Our results extend and unify various known results in literature. We also give application of proved result for four finite families of maps.

Our results extend and unify various known results in literature such as [4-14].

II. PRELIMINARIES

Definition 2.1 [14]. Two self maps A and S of a metric space are weakly compatible if $ASx = SAx$ for all x at which $Ax = Sx$.

Definition 2.2 [3]. Self maps A and S on a satisfies the property (E.A) if there exist a sequence $\{x_n\}$ in V such that $\lim_{n \rightarrow \infty} A(x_n) = \lim_{n \rightarrow \infty} S(x_n) = z$ for some $z \in V$.

Clearly, both compatible and noncompatible pairs enjoy property (E.A).

Definition 2.3 [12]. Two pairs of self maps (A, S) and (B, T) on a satisfy common property (E.A) if there exists two sequences $\{x_n\}$ and $\{y_n\}$ in V such that $\lim_{n \rightarrow \infty} A(x_n) = \lim_{n \rightarrow \infty} S(x_n) = \lim_{n \rightarrow \infty} T(y_n) = \lim_{n \rightarrow \infty} B(y_n) = p$

for some $p \in V$.

Definition 2.4 [13]. Two pairs of self maps (A, S) and (B, T) on a satisfy the $(JCLR_{ST})$ property (with respect to mappings S and T) if there exist two sequences $\{x_n\}$ and $\{y_n\}$ in V such that

$$\lim_{n \rightarrow \infty} A(x_n) = \lim_{n \rightarrow \infty} S(x_n) = \lim_{n \rightarrow \infty} T(y_n) = \lim_{n \rightarrow \infty} B(y_n) = Sz = Tz$$

where $z \in V$.

Definition 2.5 [1]. Two finite families of self maps $\{A_i\}_{i=1}^m$

and $\{B_j\}_{j=1}^n$ on a set X are

pairwise commuting if

(i) $A_i A_j = A_j A_i$, $i, j \in \{1, 2, 3, \dots, m\}$,

(ii) $B_i B_j = B_j B_i$, $i, j \in \{1, 2, 3, \dots, n\}$,

(iii) $A_i B_j = B_j A_i$, $i \in \{1, 2, 3, \dots, m\}$, $j \in \{1, 2, 3, \dots, n\}$.

III. MAIN RESULTS

Let Φ be the set of all continuous functions $\Psi: X \rightarrow X$ satisfying $\Psi(a) < a$ for all a in X .

Theorem 3.1: Let A, B, S and T be four self maps in metric space (X, d) satisfying:

(3.1) $A(X) \subset T(X)$ and $B(X) \subset S(X)$;

(3.2) there exist $\Psi \in \Phi$ such that

$$d(Ax, By) = \Psi(M(x, y))$$

where

$$M(x, y) = \max \{d(Sx, Ty), d(Sx, Ax), d(By, Ty)\}$$

for all $x, y \in X$;

(3.3) pair (A, S) or (B, T) satisfies the property (E.A).

(3.4) range of one of the maps A, B, S or T is a closed subspace of X .

Then pairs (A, S) and (B, T) have coincidence point. Further if (A, S) and (B, T) be weakly compatible pairs of self maps of X then A, B, S and T have a unique common fixed point in X .



Proof: If the pair (B, T) satisfies the property $(E.A.)$, then there exist a sequence $\{x_n\}$ in X such that $Bx_n \rightarrow z$ and $Tx_n \rightarrow z$ for some $z \in X$ as $n \rightarrow \infty$.

Since, $B(X) \subset S(X)$, therefore, there exist a sequence $\{y_n\}$ in V such that $Bx_n = Sy_n$. Hence, $Sy_n \rightarrow z$ as $n \rightarrow \infty$.

Also, since $A(X) \subset T(X)$, there exist a sequence $\{z_n\}$ in X such that $Tx_n = Az_n$. Hence, $Az_n \rightarrow z$ as $n \rightarrow \infty$.

Suppose that $S(X)$ is a closed subspace of X . Then $z = Su$ for some $u \in X$. Therefore, $Az_n \rightarrow Su, Bx_n \rightarrow Su, Tx_n \rightarrow Su, Sy_n \rightarrow Su$ as $n \rightarrow \infty$.

First we claim that $Au = Su$. Suppose not, then by (3.2), take $x = u, y = x_n$, we get

$$d(Au, Bx_n) = \Psi(M(u, x_n))$$

As $n \rightarrow \infty$

$$d(Au, Su) = \Psi\left(\lim_{n \rightarrow \infty} M(u, x_n)\right) \dots (3.5)$$

where

$$M(u, x_n) = \max\{d(Su, Tx_n), d(Su, Au), d(Bx_n, Tx_n)\}$$

As $n \rightarrow \infty$

$$\lim_{n \rightarrow \infty} M(u, x_n) = \max\{d(Su, Su), d(Su, Au), d(Su, Su)\} = d(Su, Au)$$

(3.5) gives,

$$d(Au, Su) = \Psi(d(Au, Su)) < d(Au, Su)$$

a contradiction, hence, $Au = Su$. As A and S are weakly compatible. Therefore,

$$ASu = SAu \text{ and then } AAu = ASu = SAu = SSu.$$

On the other hand, since $A(X) \subseteq T(X)$, there exist $v \in X$ such that $Au = Tv$. We now show that, $Tv = Bv$. Suppose not, then by (3.2), take $x = u, y = v$, we have,

$$d(Au, Bv) = \Psi(M(u, v))$$

or

$$d(Tv, Bv) = \Psi(M(u, v)) \dots (3.6)$$

where

$$\begin{aligned} M(u, v) &= \max\{d(Su, Tv), d(Su, Au), d(Bv, Tv)\} \\ &= \max\{d(Tv, Tv), d(Au, Au), d(Bv, Tv)\} \\ &= d(Bv, Tv) \end{aligned}$$

Thus, (3.6) gives,

$$d(Tv, Bv) = \Psi(d(Bv, Tv)) < d(Bv, Tv)$$

a contradiction, hence, $Bv = Tv$.

As B and T are weakly compatible, therefore, $BTv = TBv$ and hence, $BTv = TBv = TTv = BBv$.

Next we claim that $AAu = Au$.

Suppose not, then by (3.2), take $x = Au, y = v$, we get

$$d(AAu, Bv) = \Psi(M(Au, v)) \dots (3.7)$$

where

$$\begin{aligned} M(Au, v) &= \max\{d(SAu, Tv), d(SAu, AAu), d(Bv, Tv)\} \\ &= \max\{d(AAu, Bv), d(AAu, AAu), d(Tv, Tv)\} \\ &= d(AAu, Bv) \end{aligned}$$

(3.7) gives,

$$d(AAu, Bv) = \Psi(d(AAu, Bv)) < d(AAu, Bv)$$

again a contradiction, hence $AAu = Au$. Therefore, $Au = AAu = SAu$ and Au is a common fixed point of A and S . Similarly, we can prove that Bv is a common fixed point of B and T . As $Au = Bv$, we conclude that Au is a common fixed point of A, B, S and T .

The proof is similar when $T(X)$ is assumed to be a closed subspace of X . The cases in which $A(X)$ or $B(X)$ is a closed subspace of X are similar to the cases in which $T(X)$ or $S(X)$ respectively, is closed since $A(X) \subseteq T(X)$ and $B(X) \subseteq S(X)$.

For uniqueness, let u and v are two common fixed points of A, B, S and T . Therefore, by definition, $Au = Bu = Tu = Su = u$ and $Av = Bv = Tv = Sv = v$. Then by (3.2), take

$x = u$ and $y = v$, we get

$$d(Au, Bv) = \Psi(M(u, v))$$

or

$$d(u, v) = \Psi(M(u, v)) \dots (3.8)$$

where

$$\begin{aligned} M(u, v) &= \max\{d(Su, Tv), d(Su, Au), d(Bv, Tv)\} \\ &= \max\{d(u, v), d(u, u), d(v, v)\} \\ &= d(u, v) \end{aligned}$$

Equation (3.8) gives,

$$d(u, v) = \Psi(d(u, v)) < d(u, v)$$

a contradiction, therefore, $u = v$. Hence A, B, S and T have a unique common fixed point in X .

Taking $B = A$ and $T = S$ in Theorem 3.1, we get following result:

Corollary 3.1: Let A and S be two self maps in metric space (X, d) such that

(3.9) there exist $\Psi \in \Phi$ and $x, y \in X$ such that



$$d(Ax, Ay) = \Psi(M(x, y)) \text{ where}$$

$$M(x, y) = \max \{d(Sx, Sy), d(Sx, Ax), d(Ay, Sy)\};$$

(3.10) pair (A, S) satisfies the property $(E.A)$

(3.11) the range of one of the maps A or S is a closed subspace of X

Then A and S have a coincidence point in X . Further if (A, S) be weakly compatible pair of self maps then A and S have a unique common fixed point in X .

As an application of Theorem 3.1, we prove a common fixed point theorem for four finite families of maps. While proving our result, we utilize Definition 2.9 which is a natural extension of commutativity condition to two finite families.

Theorem 3.4: Let $\{A_1, A_2, \dots, A_m\}$, $\{B_1, B_2, \dots, B_n\}$, $\{S_1, S_2, \dots, S_p\}$ and $\{T_1, T_2, \dots, T_q\}$ be four finite families of self maps of a metric space (X, d) such that $A = A_1 A_2 \dots A_m$, $B = B_1 B_2 \dots B_n$, $S = S_1 S_2 \dots S_p$ and $T = T_1 T_2 \dots T_q$ satisfy the condition (3.2) and

$$(3.21) \quad A(X) \subset T(X) \text{ (or } B(X) \subset S(X))$$

$$(3.22) \quad \text{the pair } (A, S) \text{ (or } (B, T)) \text{ satisfy property } (E.A).$$

Then the pairs (A, S) and (B, T) have a point of coincidence each. Moreover finite families of self maps A_i , S_k , B_r and T_t have a unique common fixed point provided that the pairs of families $(\{A_i\}, \{S_k\})$ and $(\{B_r\}, \{T_t\})$ commute pairwise for all $i = 1, 2, \dots, m$, $k = 1, 2, \dots, p$, $r = 1, 2, \dots, n$, $t = 1, 2, \dots, q$.

Proof: Since self maps A, B, S, T satisfy all the conditions of theorem 3.1, the pairs (A, S) and (B, T) have a point of coincidence. Also the pairs of families $(\{A_i\}, \{S_k\})$ and $(\{B_r\}, \{T_t\})$ commute pairwise, we first show that $AS = SA$ as

$$\begin{aligned} AS &= (A_1 A_2 \dots A_m)(S_1 S_2 \dots S_p) = (A_1 A_2 \dots A_{m-1})(A_m S_1 S_2 \dots S_p) \\ &= (A_1 A_2 \dots A_{m-1})(S_1 S_2 \dots S_p A_m) = (A_1 A_2 \dots A_{m-2})(A_{m-1} S_1 S_2 \dots S_p A_m) \\ &= (A_1 A_2 \dots A_{m-2})(S_1 S_2 \dots S_p A_{m-1} A_m) = \dots = A_1 (S_1 S_2 \dots S_p A_2 \dots A_m) \\ &= (S_1 S_2 \dots S_p)(A_1 A_2 \dots A_m) = SA. \end{aligned}$$

Similarly one can prove that $BT = TB$. And hence, obviously the pair (A, S) and (B, T) are weakly compatible. Now using Theorem 3.1, we conclude that A, S, B and T have a unique common fixed point in V , say z .

Now, one needs to prove that z remains the fixed point of all the component maps.

For this consider

$$\begin{aligned} A(A_i z) &= ((A_1 A_2 \dots A_m) A_i) z = (A_1 A_2 \dots A_{m-1})(A_m A_i) z \\ &= (A_1 A_2 \dots A_{m-1})(A_i A_m) z = (A_1 A_2 \dots A_{m-2})(A_{m-1} A_i A_m) z \\ &= (A_1 A_2 \dots A_{m-2})(A_i A_{m-1} A_m) z = \dots = A_1 (A_i A_2 \dots A_m) z \\ &= (A_1 A_i)(A_2 \dots A_m) z \\ &= (A_i A_1)(A_2 \dots A_m) z = A_i (A_1 A_2 \dots A_m) z = A_i A z = A_i z. \end{aligned}$$

Similarly, one can prove that

$$\begin{aligned} A(S_k z) &= S_k (A z) = S_k z, \quad S(S_k z) = S_k (S z) = S_k z, \\ S(A_i z) &= A_i (S z) = A_i z, \quad B(B_r z) = B_r (B z) = B_r z, \\ B(T_t z) &= T_t (B z) = T_t z, \quad T(T_t z) = T_t (T z) = T_t z \end{aligned}$$

and

$$T(B_r z) = B_r (T z) = B_r z,$$

which shows that (for all i, r, k and t) $A_i z$ and $S_k z$ are other fixed point of the pair (A, S) whereas $B_r z$ and $T_t z$ are other fixed points of the pair (B, T) . As A, B, S and T have a unique common fixed point, so, we get

$$z = A_i z = S_k z = B_r z = T_t z, \quad \text{for all } i = 1, 2, \dots, m, \quad k = 1, 2, \dots, p,$$

$$r = 1, 2, \dots, n, \quad t = 1,$$

$$2, \dots, q.$$

which shows that z is a unique common fixed point of

$$\{A_i\}_{i=1}^m, \{S_k\}_{k=1}^p, \{B_r\}_{r=1}^n \text{ and } \{T_t\}_{t=1}^q.$$

IV.CONCLUSION

Using property $(E.A)$ and its variants, we prove some common fixed point theorems for quadruple of weakly compatible self maps in metric space in this paper. Our results extend and unify various known results in literature. We also give application of proved result for four finite families of maps.

V. REFERENCE

- [1] DPRVS Rao and R. Pant, Fixed point theorems in Boolean vector spaces. *Nonlinear Anal.*, Vol. 74, pp. 5383–5387, 2011.
- [2] S. Mishra, R. Pant and V. Murali, Fixed point theorems for a class of maps in metric spaces, *Fixed Point Theory and Applications*, Vol. 47 (2012).
- [3] M. Aamri and D. El Moutawakil, Some new common fixed point theorems under strict contractive condition., *J. Math. Anal. Appl.*, Vol. 270, pp. 181–188, 2002.
- [4] C. Ghilezan, “Some fixed point theorems in Boolean Algebra”, *Publ. Inst. Math. (Beograd)*, Vol. 28, Issue 42, pp. 77–82, 1980.
- [5] J. Ali, M. Imdad and D. Bahuguna, “Common fixed point theorems in Menger spaces with common property $(E.A)$ ”, *Comput. Math. Appl.*, Vol. 60, Issue 12, pp. 3152–3159, 2010.
- [6] S. Chauhan, W. Sintunavarat, P. Kumam, “Common fixed point theorems for weakly compatible



- mappings in fuzzy metric spaces Using (JCLR) property”, *Applied Mathematics*, Vol 3, Issue 9, pp. 976-982, 2012.
- [7] G. Jungck, “Compatible mappings and common fixed points”, *Internat. Jou. Math. Math. Sci.*, Vol. 9, Issue 4, pp. 771–779, 1986.
 - [8] S. Rudeanu, “Boolean transformations with unique fixed points”, *Math Slovaca.*, Vol. 57, pp. 1-10, 2007.
 - [9] S. Manro, S. S. Bhatia and S. Kumar, “Common fixed point theorem for weakly compatible maps satisfying E.A. property in intuitionistic fuzzy metric spaces”, *Punjab University Journal of Mathematics*, Vol. 42, pp. 51-56, 2010.
 - [10] S. Manro, S. S. Bhatia and Sanjay Kumar, “Common fixed point theorem for weakly compatible maps satisfying E.A. property in intuitionistic menger spaces”, *Asian Journal of Current Engineering and Maths*, Vol. 1, Issue 1, pp. 5-8, 2012.
 - [11] S. Manro, S. S. Bhatia and S. Kumar, “Common fixed point theorem for weakly compatible maps satisfying property E.A. in intuitionistic fuzzy metric spaces using implicit relation”, *Inventi Rapid Fuzzy System*, Vol. 1, pp. 1-3, 2012.
 - [12] S. Manro, S. Kumar, S. Kumar and S. S. Bhatia, “Common fixed point theorem in intuitionistic fuzzy metric spaces using common (E.A) property and implicit relation”, *Journal of Advanced Studies in Topology*, Vol. 3, pp. 60-68, 2012.
 - [13] S. Kumar, S. S. Bhatia and S. Manro, “Common fixed point theorem for weakly compatible maps satisfying E.A property in intuitionistic fuzzy metric spaces using implicit relation”, *Global Journal of Science Frontier Research Mathematics and Decision Sciences*, Vol. 12, Issue 6, pp. 25- 32, 2012.
 - [14] S. Manro, “A common fixed point theorem for weakly compatible maps satisfying property (E.A.) in fuzzy metric spaces using strict contractive condition”, *ARPN Journal of Science and Technology*, Vol. 2, Issue 4, pp. 367-370, 2012.

IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

ABOUT IJEAST

International Journal of Engineering Applied Science and Technology (IJEAST) is a peer-reviewed, open access journal that publishes high-quality research papers in the field of Engineering, Applied Science and Technology.

IJEAST aims to provide a platform for researchers, academicians, and professionals to share their innovative ideas, research findings, and practical experiences with the global scientific community.

FOCUS AREAS

- Engineering
- Applied Science
- Technology
- Innovation & Development
- Interdisciplinary Studies



PEER REVIEWED

All submissions are rigorously peer reviewed to ensure quality.



OPEN ACCESS

Free and unrestricted access to research for all.



GLOBAL REACH

Connecting researchers and professionals worldwide.



TIMELY PUBLICATION

We ensure a swift and efficient publication process.



For more information, visit our website

www.ijeast.com



INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

✉ editor@ijeast.com

🌐 www.ijeast.com

📍 India



2455-2143