



IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY



VOLUME : 4 ISSUE : 10 Print / Issue Publication Date: 09-Apr-2020



ISSN : 2455-2143



DOI : 10.33564/IJEAST.2020.v04i10.056

Indexed In



WWW.IJEAST.COM

editor@ijeast.com



LARVICIDAL ACTIVITY OF LEAF EXTRACTS OF *AGERATUM CONYZOIDES* AND *HYPTIS SUAVEOLENS* AGAINST *AEDES AEGYPTI*

Chude, C.F., Okorie, C.J., Ayegbunam, E.S., Aghalu, U.C.
Department of Zoology, Faculty of Bioscience,
Nnamdi Azikiwe University Awka, Anambra State

Abstract — Mosquito-borne diseases are the most significant public health risks globally. The potency of ethanolic leaf extracts of Goat weed (*Ageratum conyzoides*) and Pignut weed (*Hyptis suaveolens*) as mosquito larvicides under laboratory conditions were investigated. The leaf extracts of test plants were obtained using Soxhlet apparatus. Different test concentrations of 10, 8, 6, 4, 2, and 0 mg/ml (control) were prepared from stock solution. Twenty larvae of *Aedes aegypti* were exposed to various concentrations and observations were made for the period of 24 hour. The result indicates that the mean larval mortality of *A. aegypti* was recorded highest at concentration of 10 mg/ml (4.92 ± 3.605) and (3.25 ± 2.598) followed by 8 mg/ml (4.50 ± 2.747) and (2.58 ± 2.065) for leaf extracts of *A. conyzoides* and *H. suaveolens* respectively; hence, no mortality was recorded in the control. There was significant difference ($p < 0.05$) between mortality caused by concentrations and control for both plant extracts. The mean larval mortality rate of *A. aegypti* was recorded highest at 3 hours (4.33 ± 3.597) and (3.11 ± 2.698) on exposure to *A. conyzoides* and *H. suaveolens* leaf extracts respectively. There was also significant difference ($p < 0.05$) between the mortality recorded at various time of exposure. The percentage mortality was higher in leaf extracts of *Ageratum conyzoides* than *Hyptis suaveolens* but there was no significant difference ($p > 0.05$) between the mean mortality of the two plants extracts. However, the LC_{50} value for *A. conyzoides* and *H. suaveolens* were found to be 4.30 mg/ml and 1.81 mg/ml respectively for acute 24 hours test. It was recommended that leaf extracts of both plants can be used as larvicides against these insect vectors of public health importance.

Keywords — *Aedes aegypti*, Larvicidal activity, Leaf extracts, LC_{50} value

I. INTRODUCTION

Mosquito-borne diseases are the most significant public health risks globally (Webb, 2008). World health organization, has declared mosquito “public enemy number one” because they represent a significant threat to human health and their ability

to act as vector of pathogens that cause different diseases that afflict millions of people worldwide (WHO, 1998). Mosquito alone transmit diseases to more than 700 million people annually and the disease is endemic in more than 100 countries (Jang *et al.*, 2002). Beatty *et al.* (2007) reported that fifty-five percent (55%) of the world’s population are at risk of mosquito borne diseases in 124 countries. It is estimated that at least 500 million people suffer from mosquito-borne diseases (Madhumathy *et al.*, 2007).

The medical importance of mosquitoes as vectors for the transmission of serious diseases that cause morbidity, mortality, economic loss, and social disruption can never be over emphasized. These diseases have accounted for huge economic losses, mortality, low productivity and social discrimination in many developing countries (Adeleke *et al.*, 2010) and according to Mgbemena (2010), depending on the species, mosquitoes are vectors of protozoa (e.g. *Plasmodium* spp.) that cause malaria, the nematode worms that causes filariasis (e.g. *Wuchereria bancrofti*) and a large number of arboviruses including two which are of great impact in the tropical and subtropical regions (i.e. yellow fever and dengue viruses). The species of *Anopheles*, *Culex* and *Aedes* are important vectors in the transmission of malaria, filariasis, dengue, yellow and Chikungunya fevers respectively (Prajapati *et al.*, 2005). WHO (2014), implicates *Aedes* mosquito as vectors of diseases, *Aedes aegypti* is responsible for transmission of Dengue, yellow fever, Chikungunya and Zika virus while *Aedes albopictus* are responsible for the transmission of Chikungunya, dengue and West Nile Virus.

In the absence of effective prophylactic vaccine against most of the mosquito-borne diseases, it is desirable to find compounds that can effectively control mosquitoes with minimal damage to the environment. Okigbo *et al.* (2010), stated that, one of the approaches for control of these mosquito-borne diseases is the interruption of the disease transmission, by killing or preventing mosquitoes from biting human being, Larval control (source reduction or suppression) has been identified as one of the most effective methods for the control of mosquito- borne diseases as stated by (Singh *et*

al., 2006). Experience has shown that, aerial toxicants for the eradication of this mosquito are not effective, since it is highly domesticated and many adults rest indoors in hidden places such as closets (Senthilkumar *et al.*, 2008). Mosquito control has become increasingly difficult because of the indiscriminate uses of synthetic chemical insecticides which have an adverse impact on the environment and disturb the ecological balance (Nganjiwa *et al.*, 2015).

Repeated use of these controlling agents has fostered several environmental and health concerns, including disruption of natural biological control systems, outbreaks of other insect species, widespread development of resistance and undesirable effects on non- target organisms (Govindarajan *et al.*, 2013). Due to the severe effects of synthetic pesticides, there is serious need to emphasis on natural, economical, biodegradable and environmental friendly pesticide in the control of pest especially mosquitoes and that is the essence of this research. Natural products are generally preferred because of their less harmful nature to non-target organisms and due to their innate biodegradability. Plants are considered as a rich source of bioactive chemicals (Kamaraj *et al.*, 2008) and therefore an ideal control method is thus the system that uses the natural plant products as alternative to synthetic substances, as they are effective and compatible with human, animal life and the environment (Chaithong *et al.*, 2006).

Several studies showed that plant extracts or essential oils can provide desired larvicidal properties against different species of mosquitoes without causing bad effects to humans (Salunkhe *et al.*, 2011; Kumar *et al.*, 2014; Ansori *et al.*, 2015) and this is the reason for this study.

II. MATERIALS AND METHOD

Study Area

The study was conducted at the permanent site of Nnamdi Azikiwe University and Ifite environs in Awka South Local Government Area of Anambra state, Nigeria. Awka is the capital of Anambra State and located in the lowland rain forest zone of Southern Nigeria. Awka is located between latitude 5° E and 6.25° N and longitude 7° E and 8° E (Onyido *et al.*, 2010). The environment of Nnamdi Azikiwe University (Permanent site) and Ifite comprises of unharnessed thick forest filled with diversity of living organism in which *Aedes aegypti* mosquito has been reported by Okeke *et al.* (2013) to be abundant.

Collection and Identification of Plants

Twigs of pignut weeds (*Hyptis suaveolens*) (Plate 1) and goat weed (*Ageratum conyzoides*) (Plate 2) were collected from the forest of Nnamdi Azikiwe University Awka and were taken to the herbarium section in the Department of Botany Nnamdi Azikiwe University Awka, Anambra State, for identification.



Plate 1: Pignut weed (*Hyptis suaveolens*)



Plate 2: Goat weed (*Ageratum conyzoides*)

Preparation and Extraction of Test Plants

Fresh leaves were separated from the twigs and rinsed in water. 50g of fresh leaves of *Ageratum conyzoides* were weighed using electronic balance. The Soxhlet apparatus was set up and the weighed samples were packed into the Soxhlet column to 2/3 its volume. The column was then inserted into the flask and filled with 500ml of Ethanol (solvent). The soxhlet was then placed on the heating mantle, and heat was adjusted to a temperature of about 78°C. The solvents were allowed to reflux repeatedly, until refluxing solvent was clear and free from extracts.

The procedures were repeated for pignut weed (*Hyptis suaveolens*). Both extractions lasted for (6 hrs). The extracted content was transferred into a beaker, and then subjected to rotary vacuum evaporator with water-bath until solvents were completely evaporated to get the solidified crude extracts. The crude extracts thus obtained was stored in sterilized amber



coloured bottles and maintained at 4°C in a refrigerator. These extraction processes were conducted under room temperature of (28±1°C) in the Laboratory of phytochemical section of National Arbovirus and Vector Research Centre Enugu, Nigeria.

Collections, Identification and Rearing of Aedes Mosquito

Collections of eggs and larvae were carried out at permanent site of Nnamdi Azikiwe University and Ifite environs, Anambra State. The ovitraps were set under vegetation in five strategic point, larva were also collected from the natural sites such as plastic containers, used tires and stagnant water, the equipment used includes scoop- spoons, bowls, nets and specimen bottles. The eggs were collected from ovitraps, both the eggs and the larvae were sent to the Department of Entomology and Parasitology, National Arbovirus and Vector Research Centre Enugu, Nigeria, for sorting and identification. The larvae were reared under room temperature of about 28±1°C (Ahmad, 2004).

Formulation of Test Materials (Plant Extracts)

The volume of stock solution for *Ageratum conyzoides* was obtained by weighing 1g of the crude extract and 100ml of ethanol was added to dissolve it to obtain a concentration of 10mg/ml. Serial dilutions were made from this stock solution to obtain other test concentration of 10, 8, 6, 4 and 2mg/ml. These processes were repeated for *Hyptis suaveolens*.

Bioassay of Larvae with Plant Extract

In this present experiment, treatment of larvae with ethanol extracts was conducted in accordance with WHO standard method (WHO, 1986). Twenty (20) larvae of *Aedes aegypti* were placed in transparent plastic cups of about 40 ml capacity, 15 ml of tap water was added into the cups and was replicated three times. The larvae were then exposed to various concentrations of *Ageratum conyzoides* and *Hyptis suaveolens*. Control medium was also maintained with 20 larvae in 15ml of tap water for each concentration in which 1ml of ethanol was added. Mortality rates were recorded at 3, 6, 9 and 24hr in laboratory at a room temperature of 28 ± 1°C. Each of the test samples and standard were replicated 3 times. This method was modified from Okigbo *et al.* (2010).

Data Collection and Statistical Analysis

The data on larvae mortality was first recorded after 3hours, 6 hours, 12 hours and 24hours. It was subjected to analysis of variance using SPSS computer package version 20. The sample means were separated using Duncan multiple range test. The mean percentage mortality of the larva was calculated. One way analysis of variance (ANOVA) was used to compare the mean mortality of *Aedes* larvae at

concentrations and after various time of exposure to the two plants extracts at 5 % significant level.

III. RESULT

Mean Larvicidal Mortality of *A. aegypti* exposed to Various Concentrations of *A. conyzoides* and *H. suaveolens* after 24 hours

The result shown in Table 1 indicates that the mean larval mortality of *A. aegypti* was recorded highest at (4.92±3.605) on exposure to concentration of 10mg/ml in *A. conyzoides* leaf extract followed by (4.50±2.747) on exposure to concentration 8mg/ml, but lowest in the control. The result also revealed that the larval mortality of *A. aegypti* vary in a dose dependent manner. However, the effect of *A. conyzoides* leaf extract on the larval mortality was significantly (p <0.05) different between concentrations and the control.

Table 1 also indicates that the mean larval mortality of *A. aegypti* (3.25±2.598) was recorded higher on exposure to 10 mg/ml of *H. suaveolens* leaf extract followed by those of 8 mg/ml (2.58±2.065) but lowest in the control. The result also revealed that the larval mortality of *A. aegypti* vary in a dose dependent manner. However, the effect of *H. suaveolens* leaf extract on the larval mortality was significantly (p <0.05) different between concentrations and the control.

Mean Larvicidal Mortality of *A. aegypti* exposed to *A. conyzoides* and *H. suaveolens* Leaf Extracts at Various Exposure periods (in hours)

The result shown in Table 2 indicate that the mean larval mortality of *A. aegypti* (4.33±3.597) was recorded highest at 3 hours of exposure to *A. conyzoides* leaf extract followed by 6 hrs (2.56±2.526) but lowest at 9 hours (1.22±1.215). The result also revealed that the larval mortality of *A. aegypti* vary in a time dependent manner. However, the effect of *A. conyzoides* leaf extract on the larval mortality was significantly (p<0.05) different between the various time of exposure.

Table 1: Mean larvicidal mortality of *A. aegypti* exposed to various concentrations of *A. conyzoides* and *H. suaveolens* after 24 hours

Concentrations (mg/ml)	Mean Mortality of <i>A. aegypti</i> ± SD	
	<i>A. Conyzoides</i>	<i>H. suaveolens</i>
Control	ND	ND
10	4.92±3.605 ^{cb}	3.25±2.598 ^d
8	4.50±2.747 ^c	2.58±2.065 ^{cd}
6	2.58±1.621 ^b	1.75±1.357 ^{bc}
4	1.58±0.900 ^{ab}	1.42±1.165 ^b
2	1.08±0.996 ^a	0.58±0.793 ^a

Columns with different superscripts are significantly different ($p > 0.05$)
 ND: not detected

Table 2 revealed that the mean larval mortality of *A. aegypti* (3.11 ± 2.698) was recorded higher after 3 hours of exposure to *H. suaveolens* leaf extract followed by 6 hrs (1.50 ± 1.425), but lowest at 24 hours (0.83 ± 1.098). The result also revealed that the larval mortality of *A. aegypti* vary in a time dependent manner. However, the effect of *H. suaveolens* leaf extract on the larval mortality was significantly ($p < 0.05$) different between the various time of exposure.

Table 2: Mean Larvicidal mortality of *A. aegypti* exposed to various concentration of *A. conyzoides* and *H. suaveolens* at various time of exposure

Time of exposure (hrs)	Mean Mortality of <i>A. aegypti</i> $\pm$ SD <i>A. conyzoides</i>	<i>H. suaveolens</i>
3	4.33 ± 3.597^b	3.11 ± 2.698^b
6	2.56 ± 2.526^a	1.50 ± 1.425^a
9	1.22 ± 1.215^a	0.94 ± 0.873^a
24	1.67 ± 1.645^a	0.83 ± 1.098^a

Columns sharing similar superscripts are no significantly different at $p > 0.05$

Percentage mortality of *Aedes aegypti* exposed to *A. conyzoides* and *H. suaveolens* extracts

Figure 1 revealed that *A. conyzoides* extract caused the highest percentage mortality of *Aedes aegypti* larva than *H. suaveolens*. However, there was no significant difference between the mean mortality of the two plant extracts ($p > 0.05$).




Figure 1: Percentage Mortality of *Aedes* mosquito larva exposed to two plant extracts

LC₅₀ of *A. conyzoides* and *H. suaveolens* extracts against *A. aegypti* Larva

The median lethal concentration (LC₅₀) for *A. conyzoides* and *H. suaveolens* value based on Probit analysis were however found to be 4.30mg/ml and 1.81mg/ml respectively for acute 24 hours test and is as presented in figure 2 and 3 below. Figure 2 revealed a strong, linear relationship ($R^2 = 0.980$) between the mortality of *A. aegypti* exposed to *A. conyzoides*. Figure 3 also revealed a strong, linear relationship ($R^2 = 0.634$) between the mortality of *A. aegypti* exposed to *H. suaveolens*.




Figure 2: Probit regression line for response of *A. aegypti* to ethanolic extract of *A. conyzoides*




Figure 3: Probit regression line for response of *A. aegypti* to ethanolic extract of *H. suaveolens*

IV. DISCUSSION

The result of this study showed that ethanol leaf extract of *Ageratum conyzoides* possesses larvicidal activity against *Aedes aegypti*, since it caused larval mortality. This is in line



with the findings of Massuod *et al.* (2014). *Hyptis suaveolens* also possesses potency as larvicides against *Aedes aegypti*, since it also caused larval mortality. This is line with the result Dawet *et al.* (2016).

The result of Mean Larvicidal Mortality of *A. aegypti* exposed to various concentrations of *A. conyzoides* revealed that the larval mortality of *A. aegypti* vary in a dose dependent manner thus, there was increase in mortality as concentration were increased, with significant difference ($p < 0.05$) between mortality in concentrations and control. The larva mortality was also found to increase as concentration increases and the highest performance of *A. conyzoides* was at concentrations 15%, 20% and 50% in comparison to concentration 5% and 10%. Also, there was also increase in mortality as concentration increased, with significant difference ($P < 0.05$) between mortality across concentrations and control as in *A. conyzoides*. Also, this *H. suaveolens* result is line with the preliminary findings by Dawet *et al.* (2016) using Ethanol leaf extract of the same plant on *Anopheles* Mosquito Larvae. There was also increase in mortality as concentration increased with highest mortality recorded at 500mg/l.

The result of mean larval mortality rate shows that activity of leaf extracts of *A. conyzoides* and *H. suaveolens* on *A. aegypti* reduces as time passes, with significant difference ($p < 0.05$) between the various time of exposure. This shows that activity reduces as time passes. This is in agreement with the findings recently done by Dawet *et al.* (2016) using ethanol leaf extract of *Hyptis suaveolens* and *Chenopodium ambrosioides* on *Anopheles* mosquito larvae. The mortality was recorded highest at the first 3 hour of assay on *Hyptis suaveolens*. In Comparism, the percentage mortality was higher in leaf extracts of *Ageratum conyzoides* than *Hyptis suaveolens*. This is in agreement with previous findings of Okigbo *et al.* (2010), when *Azadirachta indica* (Juss) and *Ocimum gratissimum* (L.) was compared with *H. suaveolens*. The percentage mortality of Leaf extract of *Azadirachta indica* (Juss), and *Ocimum gratissimum* (L.) were greater than that of *H. suaveolens*. However, there was no significant difference ($P > 0.05$) between mortality of *A. aegypti* on exposure to *A. conyzoides* and *H. suaveolens*. This indicates that the two leaf extracts have similar effect on larval mortality of *A. aegypti*.

V. CONCLUSION

The study on larvicidal activity of leaf extract of *Ageratum conyzoides* and *Hyptis suaveolens* against *A. aegypti* showed that the ethanolic leaf extracts of both plants can be used as larvicides against *Aedes aegypti*. Since mosquitoes have developed resistance against currently used insecticides thereby, making effective control difficult and the adverse effects of these conventional insecticides on the environment and animals, including human could be amended by the use of

safer insecticides and larvicides of plant origin that could be used in the control of the vector that transmits these diseases.

VI. REFERENCE

- [1] Adeleke, M. A., Mafiana, C. F., Idowu, A. B., Sam-Wobo, S. O. and Idowu, O. A. (2010). Population dynamics of indoor sampled mosquitoes and their implication in disease transmission in Abeokuta, South-Western Nigeria. *Journal of Vector Borne Disease*, 47: 33-38.
- [2] Ahmed, H.H., Kamal, I., O.H. and Ramky, R.M. (2004). Studies on the molluscidal and Larvicidal properties of *Solanum nigrum* (L) leaves ethanol. *Journal of the Egyptian Society of Parasitology*, 31:839-843.
- [3] Ansori, A.N.M., Supriyadi, A.P. and Maria, V. (2015). Effectivity of Non Polar Extract Fraction from Key Lime (*Citrus aurantifolia*) and Kaffir Lime (*Citrus hystrix*) Leaves against *Aedes aegypti* Larvae. *Proceeding of the 5th Annual Basic Science International Conference*, Malang, Indonesia, pp131.
- [4] Beatty, M.E., Letson, W., Edgil, D.M. and Margolis, H. (2007). Estimating the total world population at risk for locally acquired dengue infection. *Proceedings of 56 the Annual Meeting of American Society of Tropical Medicine and Hygiene*, Philadelphia, Pennsylvania, USA, pp 1-8.
- [5] Chaithong, U., Wej, C., Kittichai, K., Atchariya, J, Pongsri, T., Dana, C., Daruna, C., Benjawan, T., and Benjawan P. (2006). Larvicidal effect of pepper plants on *Aedes aegypti* (L.) (Diptera: Culicidae). *Journal of Vector Ecology*, 31 (1): 138-144.
- [6] Dawet, A., Ikani, A. G. and Yaku, D. P. (2016). Larvicidal Effects of *Hyptis suaveolens* and *Chenopodium ambrosioides* on *Anopheles* Mosquito Larvae. *International Journal of Scientific & Engineering Research*, 7(1):456- 470.
- [7] Govindarajan, M., Yogalakshmi, K., Sivakumar, R. and Veerakumar K.(2013). Larvicidal property of Vetiver essential oil(*Vetiveria zizanioides*.L) A gainst *Culex tritaeniorhynchus* (Diptera:Culicidae). *International journal of currentscience and technology*, 2(1):1 -3.
- [8] Jang, Y.S., Kim, M.K., Ahn, Y.S and Lee, H.S. (2002). Larvicidal activity of Brazilian plant against *Aedes*



- aegypti* and *Culex pipiens* (Diptera: Culicidae). *Agricultural Chemistry and Biotechnology*, 4: 131–134.
- [9] Kamaraj, C., Rahuman, A.A., and Bagavan, A. (2008). Antifeedant and larvicidal effects of plant extracts against *Spodoptera litura* (F.), *Aedes aegypti* L. and *Culex quinquefasciatus* Say. *Parasitology Research*, 103(2): 325-331.
- [10] Kumar, S., Mishra, M., Wahab, N., and Warikoo R.(2014). Larvicidal, repellent, and irritant potential of the seed-derived essential oil of *Apium graveolens* against dengue vector, *Aedes aegypti* L. (Diptera: Culicidae). *Frontiers Public Health: Epidemiology*, 2 (147): 1-6.
- [11] Madhumathy, A.P., Aivazi, A. and Vijayan, V.A. (2007). Larvicidal efficacy of *Capsicum annum* against *Anopheles stephensi* and *Culex quinquefasciatus*. *Journal of Vector Borne Diseases*, 44(3):223-226.
- [12] Massuod , W. A. M. Raqib, S.M. A. Budiharjo and Mahajoena, E. (2014). Larvicidal potentiality of the bandotan (*Ageratum conyzoides*) leaves for controlling the three important species of mosquitoes (*Aedes aegypti*, *Culex quinquefasciatus* and *anopheles maculatus*). *IJCBS research paper*, 1: 33-43.
- [13] Mgbemena, I. C. (2010). Comparative Evaluation of Larvicidal Potentials of Three Plant Extracts on *Aedes aegypti*. *Journal of American Science*, 6(10): 435-440.
- [14] Nganjiwa, J.I., Pukuma, M.S., Philimon, J., and Ekundayo T.M. (2015). Evaluation of Larvicidal Properties of Leaves and Root Ethanolic Extracts of Some Plants Herbs against Fourth Instar Mosquito Larvae. *International Journal of Scientific and Research Publications*, 5(2): 1-3.
- [15] Okeke, J. J., Mogbo, T. C. And Nwakwusi, C. A. (2013). Comparative Study On The Abundance Of Mosquitoes In Temporary And Permanent Habitats In Awka, Anambra State. *American Journal of Life Science Researches*, 1(4): 192-199.
- [16] Okigbo, R.N., Okeke, J.J. and Madu, N.C. (2010). Larvicidal effects of *Azadirachta indica*, *Ocimum gratissimum* and *Hyptis suaveolens* against mosquito larvae. *Journal of Agricultural Technology*, 6(4): 703-719.
- [17] Onyido, A.E., Ezeani, A.C., Irikannu, K.C., Umeaneto, P.U., Egbuche, C.M., Chikezie, F.M. and Ugha , C.N. (2010). Anthropophilic Mosquito Species Prevalence in Nibo Community, Awka South Local Government Area, Anambra State, Southeastern Nigeria. *Journal of Epidemiology & Clinical Medicine*, 2(1): 14 – 20.
- [18] Prajapati, V. Tripathi, A. K Aggarwal, K. K. and Khanuja, S. P. S.(2005). “Insecticidal, Repellent and Oviposition-De- terrent Activity of Selected Essential Oils against *Anopheles stephensi*, *Aedes aegypti* and *Culex quinquefasciatus*,” *Bioresource Technology*, 96(16): 1749-1757.
- [19] Salunkhe, R.B, Patil, S.V, Patil, C.D, and Salunke, B.K.(2011). Larvicidal potential of silver nanoparticles synthesized using fungus *Cochliobolus lunatus* against *Aedes aegypti* (Linnaeus, 1762) and *Anopheles stephensi* Liston (Diptera: Culicidae). *Parasitology Research*, 109: 823–831.
- [20] Senthilkumar, A., Kannathasan, K., and Venkatesalu V. (2008). Chemical constituents and larvicidal property of the essential oil of *Blumea mollis* (D. Don) Merr. against *Culex quinquefasciatus*. *Parasitology Resource*, 103: 959-962.
- [21] Singh, S. B. Devi, W. R., Marina A, W. I. Devi, N. Swapana and C. B. Singh (2013). Ethnobotany, phytochemistry and pharmacology of *Ageratum conyzoides* Linn (Asteraceae). *Journal of Medicinal Plants Research*, 7(8):371-385.
- [22] W.H.O. (1986). *Instructions for determining the susceptibility or resistance of mosquito larvae to insecticides*. World Health Organization, Mimeograph Document, WHO/ VBC / 81.807, Geneva, Switzerland.
- [23] W.H.O. (1998). Report of the WHO informal consultation on the evaluation on the testing of insecticides, CTD/WHOPES/IC/96.1.Geneva, pp69
- [24] W.H.O. (2014). *A global brief on vector-borne diseases*. (WHO/DCO/WHO/2014.1), Geneva.
- [25] Webb, C.E. (2008). “Mosquito ecology: field sampling methods. *Australian Journal of Entomology*, 47(4):382–383.

IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

ABOUT IJEAST

International Journal of Engineering Applied Science and Technology (IJEAST) is a peer-reviewed, open access journal that publishes high-quality research papers in the field of Engineering, Applied Science and Technology.

IJEAST aims to provide a platform for researchers, academicians, and professionals to share their innovative ideas, research findings, and practical experiences with the global scientific community.

FOCUS AREAS

- Engineering
- Applied Science
- Technology
- Innovation & Development
- Interdisciplinary Studies



PEER REVIEWED

All submissions are rigorously peer reviewed to ensure quality.



OPEN ACCESS

Free and unrestricted access to research for all.



GLOBAL REACH

Connecting researchers and professionals worldwide.



TIMELY PUBLICATION

We ensure a swift and efficient publication process.



For more information, visit our website

www.ijeast.com



INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

✉ editor@ijeast.com

🌐 www.ijeast.com

📍 India



2455-2143