



IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY



VOLUME : 6 ISSUE : 5 Print / Issue Publication Date: 04-Dec-2021



ISSN : 2455-2143



DOI : 10.33564/IJEAST.2021.v06i05.054

Indexed In



WWW.IJEAST.COM

editor@ijeast.com



FAST DISSOLVING ORAL DISINTEGRATING FILMS OF VALSARTAN

Mr. Junaid Afsar Patel
Student
MES's College Of Pharmacy, Sonai

Prof. Katarnaware Ambadas B.
Department of Pharmaceutics
MES's College Of Pharmacy, Sonai

Abstract - Valsartan orodispersible tablets have been developed at 40-mg dose, with the intention of facilitating administration to patients experiencing problems with swallowing and hopefully, improving its poor oral bioavailability. Work started with selecting drug compatible excipients depending on differential scanning calorimetric analysis. A 3^3 full factorial design was adopted for the optimization of the tablets prepared by freeze-drying technique. The effects of the filler type, the binder type, and the binder concentration were studied. The different tablet formulas were characterized for their physical properties, weight variation, disintegration time, surface properties, wetting properties, and *in vitro* dissolution. Amongst the prepared 27 tablet formulas, formula number 6 (consisting of 4:6 valsartan:mannitol and 2% pectin) was selected to be tested *in vivo*. Oral bioavailability of two 40 mg valsartan orodispersible tablets was compared to the conventional commercial tablets after administration of a single dose to four healthy volunteers. Valsartan was monitored in plasma by high-performance liquid chromatography. The apparent rate of absorption of valsartan from the prepared tablets ($C_{max} = 2.879 \mu\text{g/ml}$, $t_{max} = 1.08 \text{ h}$) was significantly higher than that of the conventional tablets ($C_{max} = 1.471 \mu\text{g/ml}$, $t_{max} = 2.17 \text{ h}$), $P \leq 0.05$. The relative bioavailability calculated as the ratio of mean total area under the plasma concentration-time curve for the orodispersible tablets relative to the conventional ones was 135%. The results of the *in vivo* study revealed that valsartan orodispersible tablets would be advantageous with regards to improved patient compliance, rapid onset of action, and increase in bioavailability.

Key words: 3^3 full factorial design, freeze-drying technique, *in vivo* study, orodispersible tablets, valsartan

I. INTRODUCTION

Valsartan is an angiotensin II receptor antagonist used in the management of hypertension (1). It improves symptoms and quality of life in patients with chronic heart failure (2). Valsartan treatment had no demonstrable

negative effects on growth and development and is used safely as an antihypertensive agent in children less than 6 years old (3). Valsartan is rapidly absorbed following oral administration. It has a systemic availability of 0.25, which is reduced to about 0.15 by food. It is 95% protein bound and is mostly excreted as unchanged drug via the bile (4). It is given in doses of 40–160 mg once daily; this dosage is reduced in hepatic impairment, intravascular volume depletion, and renal impairment (5). The drug is available commercially as conventional tablets 40, 80, and 160 mg. Trials were done to formulate valsartan as a transdermal dosage form to overcome its low oral bioavailability (6).

- [1] Orodispersible tablets are those tablets dispersing upon contact with the moist mucosal surfaces of the oral cavity and quickly release their components without mastication or water before swallowing (7,8). The primary benefit of the orodispersible tablets is improved patient compliance due to ease of swallowing and no need for water. Difficulties with and resistance to tablet taking are common in all patient groups but are particularly prevalent in geriatric, pediatric, and psychiatric patients (9). Physical problems with swallowing (dysphagia) can exacerbate compliance problems and undermine treatment efficacy. Other benefits of orodispersible tablets include accuracy of dosage, rapid onset of action, and increase in bioavailability. The increased bioavailability of some orodispersible tablets compared to conventional tablets could be due to the dispersion in saliva and pregastric absorption. This pregastric absorption avoids first pass metabolism and can be a great advantage in drugs that undergo a great deal of hepatic metabolism (10–12). However, many orodispersible tablets showed nearly identical plasma concentration profiles as conventional tablets but with the major advantage of convenience (13). The present work is concerned with the formulation and characterization of valsartan orodispersible tablets for oral administration. Different drug compatible excipients were tried as fillers and binders. An *in vivo* study was conducted to assess the bioavailability of the selected formula in comparison with that of the conventional commercial tablets.



II. MATERIALS AND METHODS

Materials

Valsartan (particle size 2.91 μm) was obtained as a gift from International Drug Industries, Egypt. Mannitol was purchased from Winlab Laboratory Chemicals, UK. Spray-dried lactose, sucrose, and pregelatinized starch were a gift from Chemical Industries Development Company, Egypt. Sorbitol, xanthan gum, pectin (from citrus peel), gelatin (approximately 300 bloom), and acetonitrile (high-performance liquid chromatography (HPLC)) were obtained from Sigma, USA, β -cyclodextrin was purchased from Cerestar Inc., USA, sodium alginate (low molecular weight) was purchased from CDH Labs., India, Tareg® tablets, 40 mg, Novartis, Egypt. All other chemicals were of analytical grade and used as received.

Drug-Excipient Compatibility Study

This study was done to evaluate the compatibility of valsartan with pharmaceutical excipients of common use as fillers and binders. Differential scanning calorimetry (DSC) was used for screening. The specified samples were hermetically sealed in a flat bottomed aluminum pans and heated in the differential scanning calorimeter (Shimadzu Corporation, Japan) in an atmosphere of nitrogen, and the rate of flow was 25 ml/min. A temperature range of 0°C to 250°C was used, and the heating rate was 10°C/min.

Experiment Design

A 3^3 randomized full factorial design was used in the present study. In this design, three factors are evaluated, each at three levels, and experimental trials are performed at all the 27 possible combinations. The filler type (X_1), the binder type (X_2), and the binder concentration (X_3) were selected as independent variables. The composition of the different tablet formulas is listed in Table I. The oral disintegration time, percentage drug dissolved after 4 min (DP_4 min), dissolution rate at 4 min (DR_4 min), and dissolution efficiency at 30 min (DE_{30} min) were selected as dependent variables.

Table I
Composition and Responses for a 3^3 Factorial Design

Variables

Response values

Tablet formula	X_1	X_2	X_3	Percentage disintegrated (s)	DR_4 min ($\mu\text{g}/\text{min}$)	DP_4 min (%)	DE_{30} min (%)
1	–	–	–	75	11.2 ± 0.23	6.91 ± 0.08	27.6 ± 2.30
2	–	–	0	80	14.3 ± 0.45	4.85 ± 0.54	19.4 ± 2.09
3	–	–	1	90	17.5 ± 0.21	2.87 ± 0.23	11.4 ± 8.97
4	–	0	–	90	2.5 ± 0.14	24.99 ± 0.31	99.9 ± 9.07



Variables									Variables								
Response values									Response values								
Ta	X	X	X	Perc	Disint	DR _{4 min}	DP ₄	DE _{30 min} (	Ta	X	X	X	Perc	Disint	DR _{4 min}	DP ₄	DE _{30 min} (
ble				enta	egrati	(µg/min	min (	%)	ble				enta	egrati	(µg/min	min (	%)
t	1	2	3	ge	on	)	%)		t	1	2	3	ge	on	)	%)	
for				pus	time				for				pus	time			
mu				hed	(s)				mu				hed	(s)			
la				thro					la				thro				
				ugh									ugh				
				blis									blis				
				er									er				
				inta									inta				
				ct									ct				
							08			1	1			0.21	2.01	33	11
5	–	0	0	95	3.0±	24.99±	99.9	96.60±1.	8	–	1	0	70	4.2±	23.93±	95.7	95.16±4.
	1				0.34	1.85	9±0.	4		1				0.32	0.95	3±0.	41
							05									45	
6	–	0	1	100	4.2±	24.86±	99.4	96.45±0.	9	–	1	1	75	4.8±	24.49±	97.9	94.78±3.
	1				0.41	1.22	4±0.	08		1				0.41	0.89	6±.0	89
							07									.51	
7	–	1	–	65	3.1±	24.02±	96.0	95.90±3.	10	0	–	–	65	12.0±	24.92±	99.6	96.48±09
							9±2.									8±4.	



Variables									Variables								
Response values									Response values								
Ta	X	X	X	Perc	Disint	DR _{4 min}	DP ₄	DE _{30 min} (	Ta	X	X	X	Perc	Disint	DR _{4 min}	DP ₄	DE _{30 min} (
ble				enta	egrati	(µg/min	min (	%)	ble				enta	egrati	(µg/min	min (	%)
t	1	2	3	ge	on	)	%)		t	1	2	3	ge	on	)	%)	
for				pus	time				for				pus	time			
mu				hed	(s)				mu				hed	(s)			
la				thro					la				thro				
				ugh									ugh				
				blis									blis				
				er									er				
				inta									inta				
				ct									ct				
	1	1			0.22	2.14	71	1				1		0.11	1.65	04	38
11	0	–	0	75	13.9±	4.67±3.	18.6	46.47±1.	14	0	0	0	90	3.7±	19.99±	79.9	92.66±2.
	1				0.35	11	6±2.	22						0.24	0.96	9±1.	61
							44									17	
12	0	–	1	75	18.2±	3.64±2.	14.5	25.50±1.	15	0	0	1	95	5.0±	11.33±	45.3	68.90±4.
	1				0.22	88	6±2.	78						0.13	4.31	0±1.	38
							06									16	
13	0	0	–	90	3.4±	24.99±	99.9	96.31±3.	16	0	1	–	55	3.9±	24.94±	99.7	96.43±2.
							9±1.									5±4.	



Variables									Variables								
Response values									Response values								
Ta	X	X	X	Perc	Disint	DR _{4 min}	DP ₄	DE _{30 min} (	Ta	X	X	X	Perc	Disint	DR _{4 min}	DP ₄	DE _{30 min} (
ble				enta	egrati	(µg/min	min (	%)	ble				enta	egrati	(µg/min	min (	%)
t	1	2	3	ge	on	)	%)		t	1	2	3	ge	on	)	%)	
for				pus	time				for				pus	time			
mu				hed	(s)				mu				hed	(s)			
la				thro					la				thro				
				ugh									ugh				
				blis									blis				
				er									er				
				inta									inta				
				ct									ct				
			1		0.44	3.32	08	95			1	1		0.37	56	31	80
17	0	1	0	63	4.0±	24.99±	99.9	96.44±3.	20	1	—	0	35	9.0±	1.14±0.	4.56	13.63±1.
					0.31	2.87	4±3.	54			1			0.16	93	±1.0	20
							15									3	
18	0	1	1	69	6.1±	24.55±	98.2	96.22±4.	21	1	—	1	35	10.3±	1.17±0.	4.68	12.89±1.
					0.24	3.41	3±2.	08			1			0.39	87	±1.1	22
							57									7	
19	1	—	—	30	8.3±	6.58±1.	26.3	72.40±2.	22	1	0	—	40	1.3±	24.98±	99.9	96.59±4.
							4±4.									4±2.	

351



Variables					Response values					Variables					Response values				
Ta	X	X	X	Perc	Disint	DR _{4 min}	DP ₄	DE _{30 min} (		Ta	X	X	X	Perc	Disint	DR _{4 min}	DP ₄	DE _{30 min} (	
ble				enta	egrati	(µg/min	min (	%)		ble				enta	egrati	(µg/min	min (	%)	
t	1	2	3	ge	on	)	%)			t	1	2	3	ge	on	)	%)		
for				pus	time					for				pus	time				
mu				hed	(s)					mu				hed	(s)				
la				thro						la				thro					
				ugh										ugh					
				blis										blis					
				er										er					
				inta										inta					
				ct										ct					
Independent variables					Low		Medium	High		X ₃ = binder concentration (%)				0.5			1		2
X ₁ = filler type					Mannitol		Spray-dried lactose	So rbi tol		Transformed values				-1			0		1
X ₂ = binder type					Xanthan gum		Pectin	Ge lati n		<u>Open in a separate window</u> All values are mean of three readings ± S.D Preparation and Characterization of Valsartan Orodispersible Tablets A suspension containing valsartan together with other components of the preparation, filler (60 mg/tablet), and binder (0.5%, 1%, or 2% w/w of the total tablet weight) was prepared in distilled water using a magnetic stirrer (200 rpm). The tablet mix suspension was dispensed into tablet polyvinyl chloride (PVC) blister packs to result in a valsartan dose of 40 mg in each tablet. Tablet blister									



packs containing prepared tablets were then frozen in the freezer for 24 h. The frozen tablets were then lyophilized for 24 h using a Novalyph NL 500 Freeze Dryer (Savant Instruments, Holbrook, NY, USA).

Physical Properties

Twenty tablets of each formula were examined visually for their appearance. The tablet friability was expressed in terms of the percentage pushed through the PVC blister intact, and weight variation was described as the relative standard deviation (% R.S.D.) among each tablet formula.

Oral Disintegration Time

Three human subjects were involved in the determination of the disintegration time for each tablet formula; the time required for complete disintegration of the tablet when placed on the tongue was determined by tactile feedback using a stop watch.

In vitro Dissolution Study

The dissolution of valsartan from the prepared tablets was performed using the United States Pharmacopia XXVIII rotating basket, at a speed of 50 rpm in 400 ml Sorensen's phosphate buffer (pH 6.8) and at a temperature of $37 \pm 0.5^\circ\text{C}$. The study was conducted for 30 min, and aliquots each of 3 ml were withdrawn, at time intervals of 2 min, from the dissolution medium and replaced with an equivalent amount of the fresh medium. The samples were analyzed spectrophotometrically for valsartan content by measuring the absorbance at λ_{max} 250 nm against Sorensen's phosphate buffer pH 6.8 as a blank. Each experiment was carried out in triplicate.

For assessment and comparison, the dissolution rate of the drug during the first 4 min ($\text{DR}_{4 \text{ min}}$) was used. For this mean, the amount of drug dissolved per minute presented by each tablet was calculated by dividing the amount of drug dissolved in microgram by the corresponding time (14). Additionally, the dissolution profiles were evaluated on the basis of the dissolution efficiency parameter at 30 min ($\text{DE}_{30 \text{ min}}$) as described by Khan *et al.* (15) according to the following equation:

$$\text{DE} = \frac{\int_0^t y \cdot dt}{y_{100} t} \times 100$$

1

Where the integral in Eq. 1 is the area under the dissolution curve up to the dissolution time t , and y_{100} is the area of the rectangle described by 100% dissolution at the same time.

Wetting Test

The time required for a water soluble dye to diffuse throughout the entire tablet (at 2% w/w binder concentration) and reach its surface (wetting time) was measured by a slight modification of the procedure mentioned by Bi *et al.* (16).

Scanning Electron Microscopy

Scanning electron micrographs of the fracture plane of tablets were taken using electron microscope (Joel, 100S, Japan).

Oral Bioavailability Study

Subjects

Four male subjects aged 18 to 45 years were involved in the study. They were in good health based on medical history, vital signs (sitting systolic and diastolic blood pressure, heart rate, body temperature), and physical examination. Exclusion criteria included known hypersensitivity to any ingredient in the valsartan tablet, use of other drugs within 14 days before or during the trial, history of cardiovascular or pulmonary disorders, diabetes mellitus, or any other disease or condition that could interfere with the study results, a known tendency toward bleeding and a recent history of drug abuse.

Study Design

The study was a randomized, open-label, two-way crossover trial on healthy volunteers. Two treatments were used in a single dose with a washout period of 14 days. Treatment A consisted of two 40-mg tablets of the selected orodispersible valsartan tablets (formula number 6) taken without water. Treatment B consisted of two 40-mg tablets of the commercial conventional oral valsartan tablets (Tareg®) taken with 180 ml water. Both treatments were administered after an overnight fast of 10 h. The protocol was approved by the University Protection of Human Subjects Committee, and it complies with the Declaration of Helsinki and Tokyo for Humans.

Blood samples for pharmacokinetic analysis were obtained immediately before drug intake and at 0.25, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 4, 6, 8, and 24 h after dosing. Blood samples were collected in heparinized tubes and were centrifuged for 10 min at 3,000 rpm at room temperature within 1 h of collection. Separated plasma was aspirated and transferred into plastic tubes and were stored at -20°C until assayed.

Pharmacokinetic and Statistical Methods

Peak plasma concentration (C_{max} , microgram per milliliter), time to C_{max} (T_{max} , hour), the area under the plasma concentration–time curve from time zero to the

last quantifiable time point ($AUC_{(0-t)}$, microgram per hour per milliliter), AUC extrapolated to infinity ($AUC_{(0-\infty)}$, microgram per hour per milliliter), and $t_{1/2}$ (hour) were estimated by noncompartmental analysis using WinNonlin® software (version 1.5, scientific consulting, Inc, Cary, NC, USA).

The differences in average of data were compared by simple analysis of variance (one-way ANOVA) using the software SPSS (SPSS Inc., Chicago, USA). The significance of the difference was determined at 95% confidence limit ($\alpha = 0.05$).

Method of Assay

The assay of valsartan in plasma was performed by slight modifications for the high-performance liquid chromatography method adapted by Satana *et al.* (17). A thermo BDS hypersil C_{18} column ($5\ \mu$, 150×4.6 mm, Hypersil) and a mobile phase consisting of 0.02 M phosphate buffer (pH 3.2): acetonitrile (55:45 v/v) mixture were used. The flow rate was 1 ml/min, and the effluent was monitored at 225 nm using a SPD-10AVP,

ultraviolet-visible detector. Diclofenac sodium was used as an internal standard.


Before assay, the linearity, the precision, the selectivity, and the accuracy of the method were demonstrated.

[Go to:](#)

III. RESULTS AND DISCUSSION

Drug-Excipient Compatibility Study

The thermogram of valsartan showed a sharp characteristic peak at 107°C ($T_{\text{onset}} = 104.39^\circ\text{C}$, $T_{\text{endset}} = 116.94^\circ\text{C}$, and $\Delta H_f = -249.13\ \text{J/g}$) due to the melting of the solid drug. This peak was reserved in the thermograms of the 1:1 physical mixtures of the drug with spray-dried lactose, mannitol, sorbitol, xanthan gum, gelatin, and pectin, Fig. 1a, which confirms the compatibility of valsartan with those excipients. Other excipients showing incompatibilities with the drug (β -cyclodextrin, pregelatinized starch, sodium alginate, and sucrose) were avoided in the development of our tablets, Fig. 1b.



[Fig. 1](#)

DSC thermograms of valsartan with different excipients

Factorial Design

During the preparation of all the tablets, the drug:filler ratio was fixed to reduce the residual variation and increase the sensitivity of the experiment in detecting effects or changes in response due to the factors under investigation. Our preliminary work showed that drug loadings more than 40%

resulted in friable tablets. The choice of the levels of the X_1 and X_2 factors was based on results of the previous drug excipients compatibility study. The levels for factor X_3 were determined based on the data collected from previously published studies (18).



Preparation and Characterization of Valsartan Orodispersible Tablets

Fast-dissolving drug delivery systems can be manufactured by a variety of technologies, including direct compression, wet granulation, and freeze drying. Freeze drying was selected here because it results in porous tablets with rapid reconstitution in small amount of saliva in the mouth (18).

Physical Properties

All the prepared tablets were evaluated visually for their appearance, ease of removal of the blister, and integrity during removal, as well as weight variation. Percentage pushed through blister intact values are listed in Table I.

The filler type (factor X_1) showed the most pronounced effect on the physical characters of the tablets. Tablet formulas number 19–27, which contained sorbitol, shrank after freeze drying. They were sticky, friable, and not easily removed from blisters (their percentage pushed through blister intact values ranged from 20% to 50%). This is probably due to the hygroscopic properties of sorbitol. Even though variations in the levels of the other two factors (binder type and binder concentration) highly affected the percentage pushed through blister intact values, but all these values remained very low and unacceptable for handling and administration by patients. Contrary to that, tablets prepared using mannitol and spray-dried lactose as fillers showed smooth nonsticky surfaces with high integrity. It is clear that almost all the percentage pushed through blister intact values were acceptable (65–100%).

Studying the effect of the binder type (X_2) on the physical properties of the tablets, at each level of factors X_1 and X_3 , showed that pectin recorded the best appearance and the easiest removal and the highest percentage pushed through blister intact values. Pectin was followed by xanthan gum and then gelatin, which recorded the lowest percentage pushed through blister intact values.

Increasing the amount of the binder (factor X_2) improved the tablet appearance and integrity and increased the percentage pushed through blister intact. Increasing the binder amount increased the viscosity of the tablet mix suspension during preparation and resulted in more compact tablets.

% R.S.D. values for the mean tablet weights ranged from 1.3% to 3.7% for all the 27 formulas indicating high weight uniformity, data are not shown. This is because the used binders increased the viscosity of the tablet mix suspension during preparation, and the magnetic stirring kept all the tablet

components homogeneously suspended. The drug content obeyed the official limits, ranged from 97.43% to 101.2%.

Oral Disintegration Time

The results of the disintegration time are shown in Table I. Statistical analysis of these data was performed using the statistical software (StatView version 4.57, Abacus Concept). The analysis of variance including sum of squares, subsequent significant tests, and the calculation of the average values was also obtained using this software. The calculated F ratios exceeded their tabulated values for the three tested variables, filler type, binder type, and binder concentration, ($p \leq 0.05$). Additionally, all the interactions between the factors under investigation were found to have a significant effect on the disintegration time. The mean difference values between each two levels of the three factors exceeded the critical value calculated by Fisher's test (0.174) at $p \leq 0.05$, indicating a significant difference between the three levels of all factors on disintegration time as follows:

Tablets containing sorbitol as a filler recorded the lowest mean values for disintegration time 4.650 s (3.358), followed by mannitol 7.189 s (5.417) then spray-dried lactose 7.878 s (5.257), numbers between parenthesis represent S.D. This low value of sorbitol-based tablets was expected due to their poor integrity and high friability. Concerning the binder type (factor X_2) effect on disintegration time, xanthan gum tablets had the highest mean value 12.728 s (3.337) probably due to its ability to absorb water and swell-hindering disintegration. Pectin showed the lowest mean value 3.122 s (1.200). Increasing the binder amount (factor X_3) significantly prolonged the time required for complete disintegration of the tablets in mouth.

In vitro Dissolution

The calibration curve of valsartan in Sorensen's buffer pH 6.8 at $\lambda_{\max}$ 250 nm was linear over the range from 5 to 50 $\mu\text{g/mL}$ with the determination coefficient of 0.99967.

The initial drug dissolution rates during the first 4 min ($\text{DR}_{4 \text{ min}}$) and the extent of dissolution after 4 min ($\text{DP}_{4 \text{ min}}$) were calculated and shown in Table I. Additionally, the dissolution efficiency, based on 30 min, was calculated as a measure of both the rate and extent of drug dissolution. The statistical analysis revealed that the three factors (X_1 , X_2 and X_3), as well as the interactions between them, exhibited a highly significant effect on the three dissolution parameters ($\text{DP}_{4 \text{ min}}$, $\text{DR}_{4 \text{ min}}$, and $\text{DE}_{30 \text{ min}}$) at $p \leq 0.05$, Fig. 2.




Fig. 2

Interaction bar plot for the drug dissolution parameters showing the effect of filler type $\times$ binder type $\times$ binder concentration; error bars indicate 95% confidence interval. X.G xanthan gum, SDL spray-dried lactose

Concerning the effect of filler type on drug dissolution profile, tablets containing sorbitol showed significantly lower release rate and extent than those containing spray-dried lactose and mannitol at $p \leq 0.05$. The mean difference values between mannitol and spray-dried lactose did not exceed the critical differences calculated by Fisher's test for each of the dependant variables indicating a nonsignificant difference between the two fillers on the drug dissolution profile.

The binder type (factors X_2) and binder concentration (factors X_3) showed the most significant effect on the drug dissolution profile as evidenced by the highest

calculated F values for the two factors on $DP_{4\text{ min}}$, $DR_{4\text{ min}}$, and $DE_{30\text{ min}}$ (data are not shown). Xanthan gum (as a binder) showed the lowest release rate and extent (Fig. 3), $DE_{30\text{ min}}$ was 42.769% (27.705) in relation to 92.849% (8.711) and 95.466% (1.473) in case of pectin and gelatin, respectively. Most of the xanthan gum tablets remained intact after the *in vitro* dissolution test (30 min), although those tablets disintegrated in mouth in less than 12 s, this could be due to the mechanical force applied on tablets in mouth. Additionally, the presence of large volume of dissolution medium *in vitro* facilitated the water absorption and swelling of the tablets, the formed hydrated mass hindered the tablet disintegration and drug release from the tablet. Fisher's test for $DP_{4\text{ min}}$, $DR_{4\text{ min}}$, and $DE_{30\text{ min}}$ ($p \leq 0.05$) showed a significant difference between the three levels of factor X_2 on the drug dissolution profile.




Fig. 3

[Open in a separate window](#)

Effect of the binder type on the drug dissolution from valsartan orodispersible tablets prepared using mannitol and different binders at 2% concentration; error bars indicate S.D, $n = 3$

To further investigate the effect of the binder type (factor X_2) on the drug dissolution profile, the wetting test results were compared for the three binders. The time required for water to diffuse throughout the entire tablet and reach its surface (wetting time) was 3 (0.010), 5.2 (0.004), and 57 (0.103) s for tablets containing gelatin, pectin, and xanthan gum, respectively. These differences in the wetting time

values were easily justified by the scanning electron microscope images for a longitudinal section of the tablets (Fig. 4). Tablets containing pectin and gelatin as binders showed highly porous textures with deep wide pores. These pores formed channels, through which the dissolution medium penetrated facilitating tablet disintegration and drug diffusion (all the tablets completely dissolved in less than 6 min releasing 100% of their loadings; Fig. 4a, b). On the other hand, xanthan gum tablets showed a less porous sectional surface with narrow shallow pores (Fig. 4c).




Fig. 4

Scanning electron images of valsartan orodispersible tablets prepared using mannitol and **a** 2% gelatin, **b** 2% pectin, and **c** 2% xanthan gum

It is clear that the higher the binder amount, the lower the drug dissolution rate and extent from the tablets. Fisher's test proved significant differences between the three levels of factor X_3 on $DP_{4 \text{ min}}$, $DR_{4 \text{ min}}$, and $DE_{30 \text{ min}}$.

The above results show that, for factor X_1 (filler type), both mannitol and spray-dried lactose showed the best physical properties, acceptable disintegration time values, and the highest rate and extent of drug dissolution. Mannitol has more pleasant taste and feeling on dissolving in mouth than spray-dried lactose.

For binder type (factor X_2), pectin was superior concerning the physical properties and the disintegration time. Both pectin and gelatin had the fastest dissolution profiles.

Although the high level of factor X_3 (2% binder) showed longer disintegration time values and slower drug dissolution from the tablets comparative to the low and the medium levels, it recorded the best physical properties.

Based on the above discussion, formula number 6 (consisting of 4:6 valsartan:mannitol and 2% pectin) was selected to be tested *in vivo*. This formula fulfilled all the requirements for an orodispersible tablet. The tablets showed uniform smooth surfaces and pleasant taste, they were removed from blisters with 0% breakage, they had a disintegration time of 4.29 s, and they dissolved *in vitro* releasing 100% of their drug loading in 6 min with dissolution efficiency ($DE_{30 \text{ min}}$) of 96.45% and initial dissolution rate of 24.86 $\mu\text{g}/\text{min}$.

Oral Bioavailability Study

Validation of the Method of Assay

The calibration curve of valsartan was linear over the range from 20 to 3,000 ng/mL with the determination coefficient of 0.9997. The calibration curve had the regression equation of $y = 0.0067x + 0.087$, where y was the peak area ratio of valsartan to internal standard, and x was the concentration of valsartan. The within-run and between-run reproducibility of the method for plasma was calculated by analysis of replicates ($n = 3$) of samples containing known concentrations of 20, 400, and 3,000 ng/mL of valsartan. The precision of the method was described as % R.S.D. among each assay. The within-run % R.S.Ds were always below 7.9% and the between-run % R.S.Ds below 10.8%. The accuracy of the method was expressed as a percentage error of measured concentrations *versus* nominal concentrations according to the following equation:

$$\text{Mean relative error} = \frac{[\text{mean measured concentration} - \text{added concentration}] \times 100}{\text{nominal concentration}}$$

2

The within-run percentage mean relative error values were always below 8% and the between-run percentage mean relative error values below 10%. No interference with the internal standard and drug with plasma was found which proves the selectivity of the method.

Pharmacokinetic and Statistical Methods

The mean plasma concentration–time profiles for the prepared orodispersible tablets (treatment A) and the commercial conventional Tareg® tablets (treatment B) are shown in Fig. 5. In crossover design, each treatment appears an equal number of times in each period, the balancing of order of administration compensates for the period effect. If any extraneous variables affect the outcome differently in one period compared to another, all treatments may be affected equally the difference in means of the two treatments due to within and between treatment variability could be simply compared either by one-way ANOVA or paired t test (19). The pharmacokinetic parameters for both formulas in addition to the statistical analysis comparing the two formulas is summarized in Table II with p values and confidence intervals.




Fig. 5

Mean plasma concentration–time profiles of valsartan tablets; error bars indicate S.D, $n = 4$

Table II

Mean and 95% Confidence Intervals (in Parenthesis) for Pharmacokinetic Parameters of Valsartan Orodispersible Tablets and the Commercial Traditional Targ[®] Tablets Administered to Four Healthy Volunteers

Parameter	Orodispersible tablets	Traditional tablets	Statistical tests (p)
C_{max} (µg/ml)	2.8792 ± 0.244	1.4711 ± 0.553	0.016
AUC_{0-24} (µg h/ml)	14.7099 ± 4.4369 (2.6028–3.1556)	10.8703 ± 1.3003 (0.8448–2.0974)	0.224



Parameter	Orodispersible tablets	Traditional tablets	Statistical tests (p)
$AUC_{0-\infty} (\mu g \cdot h)$	15.2218 ± 4.7039 (9.8988–20.5447)	12.3417 ± 1.6627 (10.2445–14.0075)	0.343
$t_{max} (h)$	1.08 ± 0.38 2	2.167 ± 0.288	0.017
$t_{1/2} (h)$	4.88 ± 1.14 (3.59–6.17)	7.073 ± 2.778 (3.926–10.217)	0.275

Data are mean values $\pm$ S.D, $n = 4$

The mean maximum plasma concentration (C_{max}) of the prepared orodispersible tablets was significantly higher (nearly twofold) than the conventional tablets. The mean time to maximum plasma concentration (t_{max}) of treatment A was significantly shorter than that of treatment B. Thus, the estimates of both C_{max} and t_{max} showed an improved rate of absorption and hence bioavailability, which may be due to the rapid disintegration of the orodispersible tablets in the mouth and the dispersion in saliva.

The ratio of mean total area under the curve $AUC_{(0-24)}$ of treatment A to treatment B was 135%, which reflects higher amount of drug absorption over 24 h.

This could be due to some pregastric absorption of the drug. No statistically significant difference ($p = 0.224$) was found between the mean $AUC_{(0-24)}$ values for the two formulas, this statistically nonsignificant difference could be due to the small population size. Statistical comparison of half-life values did not indicate a significant difference between the test and the reference formulas.

Similarly, it was reported by Thurmman P.A. (20) that valsartan is rapidly absorbed with maximal plasma concentrations occurring 1–2 h after oral administration. The terminal half-life of valsartan was reported in various studies in the range of 3–7 h (21,22). The maximum plasma concentrations after single oral dose of valsartan reach 2–4 $\mu g/ml$ (23).

The above results reveal that the valsartan orodispersible tablets would be advantageous with regards to improved patient compliance, rapid onset of action, and increase in bioavailability.

[Go to:](#)

References

- Leidig M, Bambauer R, Kirchertz EJ, Szabã T, Handrock R, Leinung D, et al. Efficacy, safety and tolerability of valsartan 80 mg compared to irbesartan 150 mg in hypertensive patients on long-term hemodialysis. Clin Nephrol. 2008;69(6):425–32. [\[PubMed\]](#) [\[Google Scholar\]](#)
- Baumhäkel M, Müller U, Böhm M. Valsartan improves symptoms and quality of life in patients with chronic heart failure. MMW Fortschr Med. 2008;150(Suppl 1):48–53. [\[PubMed\]](#) [\[Google Scholar\]](#)
- Flynn JT, Meyers KE, Neto JP, De Paula Meneses R, Zurowska A, Bagga A, et al. Efficacy and safety of the angiotensin receptor blocker valsartan in children with hypertension aged 1 to 5 years. Hypertension. 2008;52(2):222–8. [\[PubMed\]](#)
- Grahame-Smith DG, Aronson JK. Oxford textbook of clinical pharmacology and drug therapy, 3rd ed. Oxford University Press; 2002. p. 487–8.
- Parfitt K. Martindale, The complete drug reference. 32nd, Pharmaceutical Press; 1999. p. 960.
- Rizwan M, Aqil M, Ahad A, Sultana Y, Ali MM. Transdermal delivery of valsartan: I. Effect of various terpenes. Drug Dev Ind Pharm. 2008;34(6):618–26. doi: 10.1080/03639040701833740. [\[PubMed\]](#) [\[CrossRef\]](#) [\[Google Scholar\]](#)
- European Pharmacopoeia; 4th ed., Suppl. 4.1. Council of Europe, Strasbourg, France; 2002. p. 2435.



8. Pierre H, Matthias S, Rosa M. Fast disintegrating oral dosage form. US Patent 6,083,531; 1998.
 9. Gohel M, Patel M, Amin A, Agrawal R, Dave R, Bariya N. Formulation design and optimization of mouth dissolve tablets of nimesulide using vacuum drying technique. AAPS PharmSciTech. 2004;5(3):e36. doi: 10.1208/pt050336. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
 10. Habib W, Khankari R, Hontz J. Fast-dissolving drug delivery systems, critical review in therapeutics. Drug Carrier Systems. 2000;17(1):61–72. [[PubMed](#)] [[Google Scholar](#)]
 11. Chang RK, Guo X, Burnside B, Couch R. Fast-dissolving tablets. Pharm Technol. 2000;24(6):52–8. [[Google Scholar](#)]
 12. Clarke A, Brewer F, Johnson ES, Kelly EA. Proceeding of the 122nd Annual Meeting of the American Neurological Association, 1997, M69.
 13. Seager H. Drug-delivery products and the Zydis fast-dissolving dosage form. J Pharm Pharmacol. 1998;50:375–82. [[PubMed](#)] [[Google Scholar](#)]
 14. Javadzadeh Y, Siahi-Shadbad MR, Barzegar-Jalali M, Nokhodchi A. Enhancement of dissolution rate of piroxicam using liquisolid compacts. I Farmaco. 2005;60(4):361–5. doi: 10.1016/j.farmac.2004.09.005. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
 15. Khan KA, Rhodes CT. Effect of compaction pressure on the dissolution efficiency of direct compression systems. Pharm Acta Helv. 1972;47:597–607. [[PubMed](#)] [[Google Scholar](#)]
 16. Bi Y, Sunada H, Yonezawa Y, Danjo K, Otsuka A, Iida K. Preparation and evaluation of a compressed tablet rapidly disintegrating in the oral cavity. Chem Pharm Bull (Tokyo) 1996;44:2121–7. [[PubMed](#)] [[Google Scholar](#)]
 17. Satana E, Altınay S, Göger NG, Özkan SA, Sentürk Z. J Pharm Biomed Anal. 2001;25(5–6):1009–13. doi: 10.1016/S0731-7085(01)00394-6. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
 18. Corveleyn S, Remon JP. Formulation and production of rapidly disintegrating tablets by lyophilisation using hydrochlorothiazide as a model drug. Int J Pharm. 1997;152:215–25. doi: 10.1016/S0378-5173(97)00092-6. [[CrossRef](#)] [[Google Scholar](#)]
 19. Bolton S, Remington, the science and practice of pharmacy, 20th edition, Hanson, GR, Lippincott Williams; 2002. p.142.
 20. Thurmann PA. A novel angiotensin type1 receptor antagonist. Expert Opin Pharmacother. 2000;1(2):337–50. doi: 10.1517/14656566.1.2.337. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
 21. Flesch G, Muller P, Lloyd P. Absolute bioavailability and pharmacokinetics of valsartan, an angiotensin II receptor antagonist, in man. Eur J Clin Pharmacol. 1997;52(2):115–20. doi: 10.1007/s002280050259. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
 22. Daneshtalab N, Lewanczuk RZ, Jamali F. High-performance liquid chromatographic analysis of angiotensin II receptor antagonist valsartan using a liquid extraction method. J Chromatogr B. 2002;766:345–9. doi: 10.1016/S0378-4347(01)00507-2. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
 23. Macek J, Klima J, Ptacek P. Rapid determination of valsartan in human plasma by protein precipitation and high performance liquid chromatography. J Chromatogr B. 2006;832:169–72. doi: 10.1016/j.jchromb.2005.12.035. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
- via direct saturation adjustment," Vision, Image and Signal Processing, IEE Proceedings -, vol. 152, pp. 561-574, 2005.
- F. Gonzalez and J. Hernandez, "A tutorial on Digital Watermarking ", In IEEE annual Carnahan conference on security technology, Spain, 1999.
- D. Kunder, "Multi-resolution Digital Watermarking Algorithms and Implications for Multimedia Signals", Ph.D. thesis, university of Toronto, Canada, 2001.
- J. Eggers, J. Su and B. Girod, "Robustness of a Blind Image Watermarking Scheme", Proc. IEEE Int. Conf. on Image Proc., Vancouver, 2000.
- Barni M., Bartolini F., Piva A., Multichannel watermarking of color images, IEEE Transaction on Circuits and Systems of Video Technology 12(3) (2002) 142-156.
- Kundur D., Hatzinakos D., Towards robust logo watermarking using multiresolution image fusion, IEEE Transactions on Multimedia 6 (2004) 185-197.
- C.S. Lu, H.Y.M Liao, "Multipurpose watermarking for image authentication and protection," IEEE Transaction on Image Processing, vol. 10, pp. 1579-1592, Oct. 2001.
- L. Ghouti, A. Bouridane, M.K. Ibrahim, and S. Boussakta, "Digital image watermarking using



balanced multiwavelets", *IEEE Trans. Signal Process.*, 2006, Vol. 54, No. 4, pp. 1519-1536.

P. Tay and J. Havlicek, "Image Watermarking Using Wavelets", in *Proceedings of the 2002 IEEE*, pp. II.258 – II.261, 2002.

P. Kumswat, Ki. Attakitmongcol and A. Striaew, "A New Approach for Optimization in Image Watermarking by Using Genetic Algorithms", *IEEE Transactions on Signal Processing*, Vol. 53, No. 12, pp. 4707-4719, December, 2005.

H. Daren, L. Jifuen, H. Jiwu, and L. Hongmei, "A DWT-Based Image Watermarking Algorithm", in *Proceedings of the IEEE International Conference on Multimedia and Expo*, pp. 429-432, 2001.

C. Hsu and J. Wu, "Multi-resolution Watermarking for Digital Images", *IEEE Transactions on Circuits and Systems- II*, Vol. 45, No. 8, pp. 1097-1101, August 1998.

R. Mehul, "Discrete Wavelet Transform Based Multiple Watermarking Scheme", in *Proceedings of the 2003 IEEE TENCON*, pp. 935-938, 2003.

IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

ABOUT IJEAST

International Journal of Engineering Applied Science and Technology (IJEAST) is a peer-reviewed, open access journal that publishes high-quality research papers in the field of Engineering, Applied Science and Technology.

IJEAST aims to provide a platform for researchers, academicians, and professionals to share their innovative ideas, research findings, and practical experiences with the global scientific community.

FOCUS AREAS

- Engineering
- Applied Science
- Technology
- Innovation & Development
- Interdisciplinary Studies



PEER REVIEWED

All submissions are rigorously peer reviewed to ensure quality.



OPEN ACCESS

Free and unrestricted access to research for all.



GLOBAL REACH

Connecting researchers and professionals worldwide.



TIMELY PUBLICATION

We ensure a swift and efficient publication process.



For more information, visit our website

www.ijeast.com



INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

✉ editor@ijeast.com

🌐 www.ijeast.com

📍 India



2455-2143