



IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY



VOLUME : 4 ISSUE : 06 Print / Issue Publication Date: 06-Dec-2019



ISSN : 2455-2143



DOI : 10.33564/IJEAST.2019.v04i06.006

Indexed In



WWW.IJEAST.COM

editor@ijeast.com



SOLUTION OF NONLINEAR PARTIAL DIFFERENTIAL EQUATIONS BY TRIPLE LAPLACE DECOMPOSITION METHOD

Adil Mousa

Department of Mathematics
Faculty of Science & Technology
Omdurman Islamic University, Khartoum, Sudan

Tarig M. Elzaki

Mathematics Department
Faculty of Sciences and Arts-Alkamil
University of Jeddah, Jeddah-Saudi Arabia

Abstract—in this work, we have discussed the solutions of partial differential equations nonlinear in Three Dimensions by Combining the Adomian decomposition method and triple Laplace transform and the examples application to solve partial differential equations in three-dimensions.

Keywords—Triple Laplace transform, Inverse Triple Laplace transform, Adomian decomposition method.

I. INTRODUCTION

The method was developed from 1970 to the 1990 by George Adomian the chair of the center for applied mathematics at the university of Georgia in (USA) is called Adomian decomposition method (ADM) is a semi analytical method for solving ordinary and partial nonlinear differential equations.[1-6]

P. S. Laplace (1749–1827) introduced the idea of the Laplace transform in 1782: [7-10] The Laplace transform denoted by the operator L is defined as

$$L[f(t); \rho] = \int_0^{\infty} e^{-\rho t} f(t) dt, \quad t > 0 \quad (1)$$

The main objective of the study is to extend the method of decomposition algorithm to convert one-dimensional Laplace Adomian decomposition method to three-dimensional Laplace Adomian decomposition method to solve nonlinear partial differential equations. The advantage of this method is Its capability to combining two powerful methods to obtain accurate solutions to nonlinear three-dimensional equations. Several examples are given to re-establish the effectiveness of this method. [11-13]

II. PROPOSED ALGORITHM

A. Theorems and Definitions of triple Laplace transform

Definition: 1 Let $f(x, y, t)$ be a function that can be expressed as convergent infinite series, and let $(x, y, t) \in R_3^+$, then, the triple Laplace Transform is denoted by:

$$L_3[f(x, y, t); (\sigma, \rho, \delta)] = \int_0^{\infty} \int_0^{\infty} \int_0^{\infty} e^{-\sigma x - \rho y - \delta t} f(x, y, t) dx dy dt \quad (2)$$

where $x, y, t > 0$ and σ, ρ, δ are Laplace variables, and

$$f(x, y, t) = \frac{1}{2\pi i} \int_{\alpha-i\infty}^{\alpha+i\infty} e^{\sigma x} \left[\frac{1}{2\pi i} \int_{\beta-i\infty}^{\beta+i\infty} e^{\rho y} \left[\frac{1}{2\pi i} \int_{\lambda-i\infty}^{\lambda+i\infty} e^{\delta t} f(\sigma, \rho, t) d\delta \right] d\rho \right] d\sigma$$

is the inverse triple Laplace transform denoted by L_3^{-1} .

Theorem: 2 let $f(t)$ be in S and let $F_n(\rho)$ denote Laplace transform of n th derivative, $f^{(n)}(t)$ of $f(t)$, then for $n \geq 1$

$$F_n(\rho) = \rho^n F(\rho) - \sum_{k=0}^{n-1} \rho^{n-1-k} f^{(k)}(0) \quad (3)$$

To obtain Laplace transform of partial derivative we use integration by parts, and then we have

$$\begin{aligned} L_3 \left[\frac{\partial f(x, y, t)}{\partial t} \right] &= \delta F(\sigma, \rho, \delta) - F(\sigma, \rho, 0), \\ L_3 \left[\frac{\partial^2 f(x, y, t)}{\partial t^2} \right] &= \delta^2 F(\sigma, \rho, \delta) - \delta F(\sigma, \rho, 0) - \frac{\partial F(\sigma, \rho, 0)}{\partial t}, \\ L_3 \left[\frac{\partial^3 f(x, y, t)}{\partial x \partial y \partial t} \right] : (\sigma, \rho, \delta) &= \sigma \rho \delta F(\sigma, \rho, \delta) + \sigma F(\sigma, 0, 0) + \\ &\quad \rho F(0, \rho, 0) + \delta F(0, 0, \delta) - \sigma \rho F(\sigma, \rho, 0) - \\ &\quad \sigma \delta F(\sigma, 0, \delta) - \rho \delta F(0, \rho, \delta) - F(0, 0, 0) \end{aligned} \quad (4)$$

B. Theorems and properties of triple Laplace transform Decomposition Method for PDEs –

In this section, we to illustrate the basic idea of this method, we consider a general non-linear no homogeneous partial differential equation

$$\frac{\partial^m u(x, y, t)}{\partial t^m} + Ru(x, y, t) + Nu(x, y, t) = g(x, y, t), \quad (5)$$



Where $m = 1, 2, 3$, with the initial conditions

$$\left. \frac{\partial^{m-1} u(x, y, t)}{\partial t^{m-1}} \right|_{t=0} = f_{m-1}(x, y), \quad m = 1, 2, 3, \quad (6)$$

where $\frac{\partial^m u(x, y, t)}{\partial t^m}$ is the partial derivative of the function $u(x, y, t)$ of order m ($m = 1, 2, 3, \dots$), R is the linear differential operator, N represents the general nonlinear differential operator, and $g(x, y, t)$ is the source term. Applying the triple Laplace transform (denoted in this paper by L_3) on both sides of Eq. (5), we get

$$L_3 \left[\frac{\partial^m u(x, y, t)}{\partial t^m} \right] + L_3 [Ru(x, y, t)] + L_3 [Nu(x, y, t)] = L_3 [g(x, y, t)], \quad (7)$$

Using the properties of Laplace transform, we obtain

$$\delta^m L_3 [u(x, y, t)] = \sum_{k=0}^{m-1} \delta^{m-1-k} \frac{\partial^k u(x, y, 0)}{\partial t^k} + L_3 [g(x, y, t)] - L_3 [Ru(x, y, t) + Nu(x, y, t)], \quad (8)$$

Where $m = 1, 2, 3$, and thus, we have

$$L_3 [u(x, y, t)] = \sum_{k=0}^{m-1} \delta^{-1-k} \frac{\partial^k u(x, y, 0)}{\partial t^k} +$$

$$\delta^{-m} L_3 [g(x, y, t)] - \delta^{-m} L_3 [Ru(x, y, t) + Nu(x, y, t)], \quad (9)$$

Operating the inverse transform on both sides of Eq. (9), we get

$$u(x, y, t) = G(x, y, t) - L_3^{-1} \left(\delta^{-m} E [Ru(x, y, t) + Nu(x, y, t)] \right), \quad (10)$$

where $G(x, y, t)$, represents the term arising from the source term and the prescribed initial conditions.

The second step in Laplace transform Decomposition Method, is that we represent the solution as an infinite series given below

$$u(x, y, t) = \sum_{n=0}^{\infty} u_n(x, y, t) \quad (11)$$

and the nonlinear term can be decomposed as

$$Nu(x, y, t) = \sum_{n=0}^{\infty} A_n, \quad (12)$$

where A_n are Adomian polynomials of $u_0, u_1, u_2, \dots, u_n$ and it can be calculated by the formula given below

$$A_n = \frac{1}{n!} \frac{\partial^n}{\partial \lambda^n} \left[N \sum_{i=0}^{\infty} \lambda^i u_i \right]_{\lambda=0}, \quad n = 0, 1, 2, 3, \dots \quad (13)$$

Substituting (11) and (12) in (10), we have

$$\sum_{n=0}^{\infty} u_n(x, y, t) = G(x, y, t) - L^{-1} \left(\delta^{-m} L \left[R \sum_{n=0}^{\infty} u_n + \sum_{n=0}^{\infty} A_n \right] \right) \quad (14)$$

On comparing both sides of the Eq. (14), we get

$$u_0(x, y, t) = G(x, y, t),$$

$$u_1(x, y, t) = -L^{-1} \left(\delta^{-m} L [Ru_0(x, y, t) + A_0] \right),$$

$$u_2(x, y, t) = -L^{-1} \left(\delta^{-m} L [Ru_1(x, y, t) + A_1] \right),$$

$$u_3(x, y, t) = -L^{-1} \left(\delta^{-m} L [Ru_2(x, y, t) + A_2] \right), \quad (15)$$

$\vdots$

In general, the recursive relation is given as

$$u_{n+1}(x, y, t) = -L^{-1} \left(\delta^{-m} L [Ru_n(x, y, t) + A_n] \right), \quad (16)$$

where $m = 1, 2, 3$, and $n \geq 0$;

Finally, we approximate the analytical solution $u(x, y, t)$ by truncated series

$$u(x, y, t) = \lim_{N \rightarrow \infty} \sum_{n=0}^N u_n(x, y, t) \quad (17)$$

III. EXPERIMENT AND RESULT

In this section, we apply triple Laplace decomposition method for PDEs to solve nonlinear partial differential equations of the three-dimensional.

Example 1: Consider nonlinear partial differential equation

$$u_t(x, y, t) + 6u(x, y, t)u_{xy}(x, y, t) + u_{xx}(x, y, t) = 0 \quad (18)$$

with initial conditions $u(x, y, 0) = xy$

Solution: Let us rewrite Equation (18) as:

$$u_t(x, y, t) = [-6u(x, y, t)u_{xy}(x, y, t) - u_{xx}(x, y, t)]$$

By taking the Laplace transform to both sides

$$L_3 [u_t(x, y, t)] = L_3 [-6u(x, y, t)u_{xy}(x, y, t) - u_{xx}(x, y, t)] \quad (19)$$

$$\text{Recall that } L_3 [u_t] = \delta F(\sigma, \rho, \delta) - F(\sigma, \rho, 0)$$

So Equation (19) becomes;

$$\delta F(\sigma, \rho, \delta) - F(\sigma, \rho, 0) = L_3 [-6u(x, y, t)u_{xy}(x, y, t) - u_{xx}(x, y, t)] \quad (20)$$

A taking the given initial conditions on Equation (20) and simplifying, we obtain;

$$T(\sigma, \rho, \delta) = \frac{1}{\sigma^2 \rho^2 \delta} - \frac{1}{\delta} L_3 [6u(x, y, t)u_{xy}(x, y, t) + u_{xx}(x, y, t)] \quad (21)$$

A taking the inverse triple Laplace transforms to Equation (21), we have;

$$u(x, y, t) = L_3^{-1} \left[\frac{1}{\sigma^2 \rho^2 \delta} \right] - L_3^{-1} \left\{ \frac{1}{\delta} L_3 [6u(x, y, t)u_{xy}(x, y, t) + u_{xx}(x, y, t)] \right\}$$

The resulting expression is

$$u(x, y, t) = xy -$$



$$L_3^{-1} \left\{ \frac{1}{\delta} L_3 \left[6u(x, y, t) u_{xy}(x, y, t) + u_{xx}(x, y, t) \right] \right\} \quad (22)$$

From Equation (22) let $u_0(x, y, t) = xy$

The recursive relation is given as:

$$u_{n+1}(x, y, t) = -L_3^{-1} \left\{ \frac{1}{\delta} L_3 \left[6A_n + \frac{\partial^2 u_n(x, y, t)}{\partial x^2} \right] \right\} \quad (23)$$

Note that A_n is the Adomian polynomial to decompose the nonlinear terms by using the relation:

$$A_n = \frac{1}{n!} \frac{d^n}{d\lambda^n} f \left[\sum_{i=0}^{\infty} \lambda^i u_i(x, y, t) \right]_{\lambda=0} \quad (24)$$

Let the nonlinear term be represented

$$f[u(x, y, t)] = u(x, y, t) \frac{\partial^2 u(x, y, t)}{\partial x \partial y} \quad (25)$$

By using Equation (25) in Equation (24), we obtain;

$$\begin{aligned} A_0 &= u_0(x, y, t) \frac{\partial^2 u_0(x, y, t)}{\partial x \partial y}, \\ A_1 &= u_1(x, y, t) \frac{\partial^2 u_0(x, y, t)}{\partial x \partial y} + u_0(x, y, t) \frac{\partial^2 u_1(x, y, t)}{\partial x \partial y}, \\ A_2 &= u_2(x, y, t) \frac{\partial^2 u_0(x, y, t)}{\partial x \partial y} + \\ &u_1(x, y, t) \frac{\partial^2 u_1(x, y, t)}{\partial x \partial y} + u_0(x, y, t) \frac{\partial^2 u_2(x, y, t)}{\partial x \partial y}, \dots \end{aligned}$$

From Equation (23)

When $n=0$, we have

$$u_1(x, y, t) = -L_3^{-1} \left\{ \frac{1}{\delta} L_3 \left[6A_0 + \frac{\partial^2 u_0(x, y, t)}{\partial x^2} \right] \right\}$$

is computed as: $A_0 = xy$

$$u_1(x, y, t) = -L_3^{-1} \left\{ \frac{1}{\delta} L_3 \left[6xy + \frac{\partial^2}{\partial x^2} [xy] \right] \right\}$$

By simplifying $u_1(x, y, t) = -6xyt$

When $n = 1$, we have;

$$u_2(x, y, t) = -L_3^{-1} \left\{ \frac{1}{\delta} L_3 \left[6A_1 + \frac{\partial^2 u_1(x, y, t)}{\partial x^2} \right] \right\}$$

is computed as: $A_1 = -12xyt$

$$u_2(x, y, t) = -L_3^{-1} \left\{ \frac{1}{\delta} L_3 \left[-72xyt + \frac{\partial^2}{\partial x^2} [-6xyt] \right] \right\}$$

By simplifying $u_2(x, y, t) = 36xyt^2$

The approximate series solution is

$$u(x, y, t) = xy - 6xyt + 36xyt^2 - 216xyt^3 + \dots$$

This can be written as

$$u(x, y, t) = xy \left[1 - 6t + [6t]^2 - [6t]^3 + \dots \right]$$

By using Taylor's series, the closed form solution will be as follows

$$u(x, y, t) = \frac{xy}{1+6t}$$

Example 2: Consider nonlinear partial differential equation

$$u_y(x, y, t) u_t(x, y, t) - u_{xx}(x, y, t) = u(x, y, t) \quad (26)$$

with initial conditions

$$u(0, y, t) = yt, \quad u_x(0, y, t) = -1$$

Solution: Let us rewrite Equation (26) as:

$$u_{xx}(x, y, t) = u_y(x, y, t) u_t(x, y, t) - u(x, y, t)$$

By taking the triple Laplace transform to both sides

$$L_3[u_{xx}(x, y, t)] = L_3[u_y(x, y, t) u_t(x, y, t) - u(x, y, t)] \quad (27)$$

Recall that

$$L_3[u_{xx}(x, y, t)] = \sigma^2 F(\sigma, \rho, \delta) - \sigma F(0, \rho, \delta) - \frac{\partial F(0, \rho, \delta)}{\partial x}$$

So Equation (27) becomes;

$$\sigma^2 F(\sigma, \rho, \delta) - \sigma F(0, \rho, \delta) - \frac{\partial F(0, \rho, \delta)}{\partial x} =$$

$$L_3[u_y(x, y, t) u_t(x, y, t) - u(x, y, t)]$$

(28)

taking the given initial conditions on Equation (28) and simplifying, we obtain;

$$\begin{aligned} F(\sigma, \rho, \delta) &= \frac{1}{\sigma \rho^2 \delta^2} - \frac{1}{\sigma^2 \rho \delta} - \\ &\frac{1}{\sigma^2} L_3[u(x, y, t) - u_y(x, y, t) u_t(x, y, t)] \end{aligned} \quad (29)$$

A taking the inverse triple Laplace transform to Eq. (29), we have;

$$\begin{aligned} u(x, y, t) &= L_3^{-1} \left[\frac{1}{\sigma \rho^2 \delta^2} - \frac{1}{\sigma^2 \rho \delta} \right] - \\ &L_3^{-1} \left\{ \frac{1}{\sigma^2} L_3[u(x, y, t) - u_y(x, y, t) u_t(x, y, t)] \right\} \end{aligned}$$

The resulting expression is

$$u(x, y, t) = yt - x -$$

$$L_3^{-1} \left\{ \frac{1}{\sigma^2} L_3[u(x, y, t) - u_y(x, y, t) u_t(x, y, t)] \right\} \quad (30)$$

From Equation (30) let $u_0(x, y, t) = yt - x$

The recursive relation is given as:

$$u_{n+1}(x, y, t) = -L_3^{-1} \left\{ \frac{1}{\sigma^2} L_3 \left[A_n - \frac{\partial u_n(x, y, t)}{\partial y} \frac{\partial u_n(x, y, t)}{\partial t} \right] \right\} \quad (31)$$



Note that A_n is the Adomian polynomial to decompose the nonlinear terms by using the relation:

$$A_n = \frac{1}{n!} \frac{d^n}{d\lambda^n} f \left[\sum_{i=0}^{\infty} \lambda^i u_i(x, y, t) \right]_{\lambda=0} \quad (32)$$

Let the nonlinear term be represented

$$f[u(x, y, t)] = u(x, y, t) \quad (33)$$

By using Equation (33) in Equation (32), we obtain;

$$A_0 = u_0(x, y, t),$$

$$A_1 = u_1(x, y, t) + u_0(x, y, t),$$

$$A_2 = u_2(x, y, t) + u_1(x, y, t) + u_0(x, y, t), \dots$$

From equation (31), when $n=0$, we have:

$$u_1(x, y, t) = -L_3^{-1} \left\{ \frac{1}{\sigma^2} L_3 \left[A_0 - \frac{\partial u_0(x, y, t)}{\partial y} \frac{\partial u_0(x, y, t)}{\partial t} \right] \right\}$$

is computed as: $A_0 = yt - x$

$$u_1(x, y, t) = -L_3^{-1} \left\{ \frac{1}{\sigma^2} L_3 \left[[yt - x] - \frac{\partial [yt - x]}{\partial y} \frac{\partial [yt - x]}{\partial t} \right] \right\}$$

$$\text{By simplifying } u_1(x, y, t) = \frac{x^3}{3!}$$

The approximate series solution is

$$u(x, y, t) = yt - x + \frac{x^3}{3!} - \frac{x^5}{5!} - \frac{1}{2} x^2 yt + \frac{x^3}{3!} \dots$$

This can be written as

$$u(x, y, t) = yt - \left[x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots \right]$$

By using Taylor's series, the closed form solution will be as follows

$$u(x, y, t) = yt - \sin x$$

IV. CONCLUSION

This work discussed the definition of the triple Laplace transform that was applied. Some important theorems and properties have been presented for this relatively new transformation to find solutions for partial differential equations in three dimensions under the initial conditions, the triple Laplace transform study succeeded in achieving solutions.

V. REFERENCE

- [1] Abdul Majid Wazwaz, Partial Differential Equations and Solitary Waves Theory, Higher Education Press Beijing and Springer - Verlag Berlin Heidelberg (2009).
- [2] Moghadam MM, Saeedi H Application of differential transforms for solving the Volterra integro-partial differential equations. Iranian Journal of Science and Technology, Transaction A 34(A1):59 (2010).

- [3] Garci'a-Olivares, A., Analytic solution of partial differential equations with Adomian's decomposition, Kybernetes, 32 (3) 354-368 (2003).
- [4] Tarig M. Elzaki, Adil Mousa On the convergence of triple Elzaki transform SN Applied Sciences 1:275 <https://doi.org/10.1007/s42452-019-0257-2> (2019)
- [5] H. Eltayeb, A. Kilicman and S. Mesloub, Modified Laplace decomposition method for solving system of equations Emden-Fowler type, Journal of Computational and Theoretical Nanoscience, 12, 5297-5301, (2015).
- [6] S.A. Khuri, A Laplace decomposition algorithm applied to a class of nonlinear differential equations, J.Math. Annl. Appl., 4, 141-155, (2001).
- [7] A. Khuri, A Laplace decomposition algorithm applied to a class of nonlinear differential equations, J. Math. Annl. Appl., 4, 141-155, (2001).
- [8] Abdon Atangana A Note on the Triple Laplace Transform and Its Applications to Some Kind of Third-Order Differential Equation Abstract and Applied Analysis, Article ID 769102, 10 pages Volume (2013).
- [9] M. Khalid, M. Sultana, F. Zaidi and U. Arshad, An Elzaki Transform Decomposition Algorithm Applied to a Class of Non-Linear Differential Equations, J. Natural Sci. Res., 5, 48-56, (2015).
- [10] Trig M. Elzaki, Salih M. Elzaki, On connection between Laplace transform and Elzaki Transform, Advances in Theoretical and Applied Mathematics, ISSN0973-4554 volume 6 Number1, pp. 1-11, (211).
- [11] M. M. Hosseini and H. Nasabzadeh, On the convergence of Adomian decomposition method, Appl. Math. Comput., 182, 536-543, (2006).
- [12] D. Kumar, J. Singh and S. Rathore, Sumudu Decomposition Method for Nonlinear Equations, Int. Math. For., 7, 515 - 521.57-64, (2012).
- [13] J.S. Duan and R. Rach, A new modification of the Adomian decomposition method for solving boundary value problems for higher order nonlinear differential equations, Appl. Math. Comput, 4090-4118, 218 (2011).

IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

ABOUT IJEAST

International Journal of Engineering Applied Science and Technology (IJEAST) is a peer-reviewed, open access journal that publishes high-quality research papers in the field of Engineering, Applied Science and Technology.

IJEAST aims to provide a platform for researchers, academicians, and professionals to share their innovative ideas, research findings, and practical experiences with the global scientific community.

FOCUS AREAS

- Engineering
- Applied Science
- Technology
- Innovation & Development
- Interdisciplinary Studies



PEER REVIEWED

All submissions are rigorously peer reviewed to ensure quality.



OPEN ACCESS

Free and unrestricted access to research for all.



GLOBAL REACH

Connecting researchers and professionals worldwide.



TIMELY PUBLICATION

We ensure a swift and efficient publication process.



For more information, visit our website

www.ijeast.com



INTERNATIONAL JOURNAL

OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY



editor@ijeast.com



www.ijeast.com



India



2455-2143