



IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY



VOLUME : 4 ISSUE : 11 Print / Issue Publication Date: 10-May-2020



ISSN : 2455-2143



DOI : 10.33564/IJEAST.2020.v04i11.082

Indexed In



WWW.IJEAST.COM

editor@ijeast.com



FOLLICLE STIMULATION HORMONE (FSH) AFFECTED BY ORAL ADMINISTRATION OF THE ENVIRONMENTAL CONTAMINANT ATRAZINE HERBICIDE BY USING *WISTAR RATS* AS EXPERIMENTAL ANIMALS, AND ITS CONFLICTS ON HUMAN HEALTH

Daradkeh S.M, Abdelgadir M.I

Nursing Department,

Alghad International Colleges for Applied Health Sciences,
Qassim, Saudi Arabia

Awad Eljeed B.A.

Chemistry Department, Faculty of Science,
Bakht Er-Ruda University, White Nile State, Sudan

Daradkeh M.S

Faculty of Veterinary Medicine,
Jordan University of Science and Technology. Jordan

Murwan K. Sabahelkhier

Department of Biochemistry and molecular biology,
Faculty of Science and Technology, El Neelain University, Khartoum, Sudan

Abstract— Twenty *Wistar rats* were divided into four groups; three groups served as test groups which were orally administered by different doses of the herbicide: (27.3, 38.5, and 42.0 mg/kg b.w), while the remaining group was put as control. Animals were freely accessed to water tab, and basal diet. Metal cages were interval cleaned. Cages were kept in 15 C degree. Decreased concentrations ($p \leq 0.05$) of FSH compared to control were observed through the course of the experiment, more precisely at the 30th day. This information could affect the endocrine disruptor-screening program that is focused on hormone mimetic compounds and regulatory systems in order to better protect the reproductive health of women from ATR exposure [5]. Some sexually selected traits concerning many vertebrate species, especially in males, in which sex hormones interpresion was established [6].

Keywords— Atrazine, Endocrine, infertile, environmental pollution, expression, steroidogenesis

I. INTRODUCTION

It has been estimated that 72.4 million women are currently infertile [2]. An ongoing and rapid increase in reproductive diseases and a decline in reproductive functions have been observed in recent years, almost due to environmental pollution by chemicals [17]. Endocrine-disrupting chemicals impacts on cellular and molecular events in the ovary and their role in ovarian pathophysiology could provide important insights into the potential reproductive harm in humans. Atrazine, is the most heavily applied agricultural pesticide for crop production worldwide. Atrazine is banned in European Union countries, while the maximal contaminant level of 0.003 mg/L in drinking water is currently under review [18]. An increased concern that agricultural runoff of this chemical can affect non-target animal species. The actual ATR levels in the environment can exceed maximal permissible concentrations MPCs, sometimes reaching more than 0.02 mg/L [15]. ATR exposure in humans is primarily related to ingestion of contaminated drinking water and from dermal contact or ingestion following agricultural and lawn

applications. Atrazine is toxicologically important, primarily because it acts as an endocrine disruptor and has a potential to interfere with steroidogenesis. Among the steroid-disrupting outcomes, the effect of ATR on aromatase (CYP19A1), an enzyme involved in estradiol synthesis [16]. Premature luteinization is a possible cause of infertility in women. It is currently unknown whether environmental chemicals can induce changes associated with premature luteinization. Using rat granulosa cells (GC) in vitro, it was demonstrated that exposure to atrazine (ATR), a widely used herbicide, causes GC phenotype that resembles that of human premature luteinization. At the end of the 48-h stimulation with FSH, ATR-exposed GC showed, higher levels of progesterone, overexpression of luteal markers, and an increase in progesterone: estradiol ratio above 1[11]. It was concluded that disorders in reproductive functions in many mammals was due to atrazine exposure. Treatment of human cumulus granulosa cells with 20 μ M ATR for 48 h resulted in lower FSH-stimulated estradiol and progesterone production. ATR reduced mRNA levels of aromatase (CYP19A1), steroidogenic acute regulatory protein (STAR) and luteinizing hormone/choriogonadotropin receptor (LHCGR). It is suggested that ATR alters steroidogenesis and ovulatory process in human cumulus granulosa cells jeopardizing female reproduction. Women who are exposed to this pesticide are more likely to experience side effects of the FSH action during regular menstrual cycling or in vitro fertilization. The concept of endocrine disruptors as hormone sensitizers has been recognized and represents a new mode of action different from the hormone mimetics. This information could affect the endocrine disruptor-screening program that is focused on hormone mimetic compounds and regulatory systems in order to better protect the reproductive health of women from ATR exposure [5]. Some sexually selected traits concerning many vertebrate species, especially in males, in which sex hormones interpresion was established [6]. Disruption of the expression or perception of such traits can influence mate choice and evolutionary patterns [14]. Increased pollution is becoming an important factor in such disruptions [13], and is hence instrumental in shaping evolutionary trajectories. Endocrine disrupting compounds (EDCs) are concerned as pollutants, which interfere with proper hormonal functioning. Many of them have anthropogenic sources such as pesticides, industrial effluents, pulp mill effluents, plastics and sewage. Significant routes of exposure include direct exposures from living in contaminated soil or water, as well as indirect exposures through eating contaminated prey [9]. All aspects of the reproductive system can be affected by EDCs, including gonadal formation, production of hormones and gametes, sex determination [1].

II. EXPERIMENT AND RESULT

Atrazine with different concentrations was prepared in a suitable carrier. Blood samples were collected interval every

ten days for forty days, then centrifuged and kept at 5 C degree, till analysis according to the method described by [4].




Fig.1. Mean values of FSH in IU/L vs Dosage mg/kg bw

Table 1. Mean difference of FSH concentration IU/L, Weight in mg/kg bw, and Time in days

	Test Value = 0					
	t	Df	Sig (2-tail ed)	Mean Difference	95% Confidence Interval of the Difference	
					Lower	Upper
FSH IU/L	3.663	15	.00 2	1.447500	.6053 0	2.289 70
mg/kg bw	6.102	15	.00 0	26.05000 0	16.95 056	35.14 944
Days	8.660	15	.00 0	25.000	18.85	31.15

This study resulted in decreased FSH concentrations in accordance to time intervals (10th, 20th, 30th, and 40th day of the experiment), as shown in Figure No.1 and Table No.1. Study results agree with [7] who demonstrated that ATR decreases the FSH-stimulated aromatase expression and consequently diminishes estradiol production. Study results are also confirmed by [10] who has shown that ATR decreases estradiol production and Cyp19a1 and Lhr expression in the FSH-stimulated GC. Study results are confirmed by [3], who stated that, in the luteal-follicle phase transition period the serum levels of progesterone and estrogen (primarily estradiol) decrease and no longer suppress the release of FSH.



III. CONCLUSION

Authors highly recommend restricting chemical weed control, and prefer usage of integrated pest management, which can be defined as an ecosystem-based strategy that focuses on long-term prevention of pests or their damage through a combination of techniques such as biological control, habitat manipulation, modification of cultural practices, and use of resistant varieties.

IV. ACKNOWLEDGEMENTS

Authors gratefully thank the family of Meck Nimir Research Center, Khartoum, Sudan, for engaging experimental rats, & Family of Khartoum hospital for sample analysis. & Dr. Eljack M.Elamin., (the Head administrator of Zagross company for import/ export of agricultural requirements), for valuable providing atrazine reagent, & Dr. Mohamed H. Bani Khaled for great financial participation and encouragement.

V. REFERENCE

- [1]. Basrur PK. Disrupted sex differentiation and feminization of man and domestic animals. *Environ Res*, Vol. 100, pp. 18–38, 2006.
- [2]. Boivin J., Bunting L., Collins JA., and Nygren KG. International estimates of infertility prevalence and treatment-seeking: potential need and demand for infertility medical care. *Hum Reprod*, Vol. 22, pp. 1506-1512, 2007.
- [3]. Fowler PA., Sorsa-Leslie T., Harris W., and Mason HD. ("Ovarian gonadotrophin surge-attenuating factor (GnSAF): where are we after 20 years of research?" (PDF). *Reproduction*, Vol.126, No. 6, pp. 689–99, 2003.
- [4]. Hinkle J., and Cheever K. Brunner and Suddarth's. *Handbook of Laboratory and Diagnostic Tests*. 2nd Ed, Kindle. Philadelphia: Wolters Kluwer Health, Lippincott Williams & Wilkins; Follicle Stimulating Hormone (FSH) Serum, pp. 306–7. 2014.
- [5]. Jansen MS., Nagel SC., Miranda PJ., Lobenhofer EK., Afshari CA., and McDonnell DP. Short-chain fatty acids enhance nuclear receptor activity through mitogen-activated protein kinase activation and histone deacetylase inhibition. *Proc Natl Acad Sci U S A*, Vol. 101, pp. 719–7204, 2004.
- [6]. Jayasooriya AP., Weisinger RS., Weisinger HS., Mathai ML., and Sinclair AJ. Attraction to orange: sexiness, not gluttony. *Science*, Vol. 296, pp. 847–848, 2002.
- [7]. Kristina P., Dragana S., Svetlana F., Jelena H., Branka G., Sonja K., and Nebojsa A. Atrazine Enhances Progesterone Production Through Activation of Multiple Signaling Pathways in FSH-Stimulated Rat Granulosa Cells: Evidence for Premature Luteinization1. *Biology of Reproduction*, Vol. 91, No. 5, pp. 1-10, 2014.
- [8]. Manzon G., and Denver RJ. Regulation of pituitary thyrotropin gene expression during *Xenopus* metamorphosis: negative feedback is functional throughout metamorphosis. *J Endocrinol*, Vol.182, pp. 273–285, 2004.
- [9]. Markman S., Guschina IA., Barnsley S., Buchanan KL., and Pascoe D. Endocrine disrupting chemicals accumulate in earthworms exposed to sewage effluent. *Chemosphere*, Vol. 70, pp. 119–125, 2007.
- [10]. Pogrmic-Majkic FS., Samardzija KD., Glisic B., Kaisarevic S., Kovacevic R., and Andric N. Involvement of ERK1/2 signaling pathway in atrazine action on FSH-stimulated LHR and CYP19A1 expression in rat granulosa cells. *Toxicol Appl Pharmacol*, Vol. 270, pp. 1–8, 2013.
- [11]. Pogrmic-Majkic K., Samardzija D., Fa S., Hrubik J., Glisic B., Kaisarevic S., and Andric N. Atrazine enhances progesterone production through activation of multiple signaling pathways in FSH-stimulated rat granulosa cells: evidence for premature luteinization. *Biol Reprod*, Vol. 91, No. 5, pp. 124, 2014.
- [12]. Pogrmic-Majkic K., Samardzija D., Stojkov-Mimic N., Vukosavljevic J., Trninic-Pjevic A., Kopitovic V., and Andric N. Atrazine suppresses FSH-induced steroidogenesis and LH-dependent expression of ovulatory genes through PDE-cAMP signaling pathway in human cumulus granulosa cells. *Mol Cell Endocrinol*, Vol. 5, No. 461, pp. 79-88, 2018.
- [13]. Seehausen O., Van Alphen JM., and Witte F. Cichlid fish diversity threatened by eutrophication that curbs sexual selection. *Science*, Vol. 277, pp. 1808–1811, 1997.
- [14]. Seehausen O., and Van Alphen JM. The effect of male coloration on female mate choice in closely related Lake Victoria cichlids (*Haplochromis nyererei* complex). *Behav Ecol Sociobiol*, Vol. 42, pp. 1–8, 1998.
- [15]. Solomon KR., Baker DB., Richards RP., Dixon DR., Klaine SJ., LaPoint TW., Kendall RJ., Weisskopf CP., Giddings JM., Giesy JP., Hall LW., and Williams WM. Ecological risk assessment of atrazine in North American surface waters. *Environ Toxicol Chem*, Vol.15, pp. 31–74, 1996.
- [16]. Tinfo NS., Hotchkiss MG., Buckalew AR., Zorrilla LM., Cooper RL., and Laws SC. Understanding the effects of atrazine on steroidogenesis in rat granulosa and H295R adrenal cortical carcinoma cells. *Reprod Toxicol*, Vol. 3, pp. 184–193, 2011.
- [17]. Wang W., and Flaws JA. Endocrine-disrupting chemicals in ovarian function: effects on steroidogenesis, metabolism and nuclear receptor signaling. *Reproductio*, Vol.142, pp. 633-646, 2011.
- [18]. Weber GJ., Sepulveda MS., Peterson SM., Lewis SS., and Freeman JL. Transcriptome alterations following developmental atrazine exposure in zebrafish are associated with disruption of neuroendocrine and reproductive system function, cell cycle, and carcinogenesis. *Toxicol Sci*, Vol. 132, pp. 458-466, 2013.

IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

ABOUT IJEAST

International Journal of Engineering Applied Science and Technology (IJEAST) is a peer-reviewed, open access journal that publishes high-quality research papers in the field of Engineering, Applied Science and Technology.

IJEAST aims to provide a platform for researchers, academicians, and professionals to share their innovative ideas, research findings, and practical experiences with the global scientific community.

FOCUS AREAS

- Engineering
- Applied Science
- Technology
- Innovation & Development
- Interdisciplinary Studies



PEER REVIEWED

All submissions are rigorously peer reviewed to ensure quality.



OPEN ACCESS

Free and unrestricted access to research for all.



GLOBAL REACH

Connecting researchers and professionals worldwide.



TIMELY PUBLICATION

We ensure a swift and efficient publication process.



For more information, visit our website
www.ijeast.com



INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

✉ editor@ijeast.com

🌐 www.ijeast.com

📍 India



2455-2143