



A REVIEW OF *TRIDAX PROCUMBENS* AND ITS ACTIVITY

Sabarinath.K, Sandhiya.S, Ishwarya.R, Logeshwaran.V, Kousalya.N

Postgraduate student

Department of Biotechnology, Dr. N.G.P. Arts and Science College (*Autonomous*)
Coimbatore-48

Arun. P

Assistant professor

Department of Biotechnology, Dr. N.G.P. Arts and Science College (*Autonomous*)
Coimbatore-48

Abstract— *Tridax procumbens* (*T. procumbens*) is also known as coat button or tridax daisy. It is the widespread weed and also a pest plant in tropical and subtropical. *T. procumbens* was used as a traditional medicine in wound healing, antifungal, antibacterial, insect repellent all over the world. The raw leave extract of *T. procumbens* is used as a best for wound healing as a Ayurvedic medicine in India.

Keywords— *Tridax procumbens*, Ayurvedic medicine

I. INTRODUCTION

Tridax procumbens (*T. procumbens*) is belong to Asteraceae family. It's a annual and perianal weed, widespread throughout India. It has bisexual flower with white headed flower and the whole plant has the activity of wound healing, antifungal, antibacterial, insect repellent, and antimalarial. The main aim of the review is to enrich the knowledge about *Tridax procumbens* and its activity.

SCIENTIFIC CLASSIFICATION

Kingdom : Plantae
Order : Asterales
Family : Asteraceae
Genus : Tridax
Species : procumbens

II. TRADITIONAL USE

The whole plant *T. procumbens* is used as a raw medicine for protozoal infection, chronic ulcer, diabetes, insect repellent, wound healing, and inflammation, jaundice, anti-malarial, antibacterial, and anti-fungal. Flavonoids which found in the whole tridax plant act as a best anticoagulants and wound healing.

III. ANTICOAGULATION ACTIVITY

200 mg/ μ g of *T. procumbens* is injected to rabbit, result in prolongation of clottind time by reduce the production of heparin.

IV. WOUND HEALING ACTIVITY

Extraction of *T. procumbens* increased the lysyl oxidase activity, protein content, and breaking strength which helps in promoting wound healing. It increased the interaction between epidermal and dermal cells.

The tridax extract also increased the glycosaminoglycan level as it increased the protein and nucleic acid content.

V. ANTIOXIDANT ACTIVITY

Methanol was used as a solvent to extract the aerial part of the plant and subjected to DPPH assay. The extraction on the solvents Ethyl acetate and n-butanol showed better antioxidant result when compared to the standard Ascorbic acid.

The ethanolic extract of tridax shows 96.70% of antioxidant where standard gallic acid and ascorbic acid shows 92.92% and 94.81% respectively.

VI. INSECTICIDAL ACTIVITY

The oil of tridax is used as strong insect repellent against house flies, mosquito's larvae, cockroaches, and *Dysder cussimile*. It is observed that tridax is neither attacked by insects and cattles.

VII. ANTIMALARIAL ACTIVITY

The oil of tridax shows relatively high antimalarial activity against *Anopheles stephensi* when compared to their two plants *I. cairica*, *M. charantia*. Clear dosage relationship



was established in all essential oils, with highest concentration of 6%.

VIII. HAIR GROWTH PROMOTING ACTIVITY

The ethanolic extract of aerial part of the plant was administered to albino rat as 10% ointment and also as orally 100g/kg for about 20 days and the both treat showed good results. The petrol fraction of the extract shows a more effective when compared to other fraction extract.

IX. ANTIPROTOZOAL ACTIVITY

Extract of 13 plant was subjected against protozoal infection, were evaluated *in vitro* against epimastigotes and tripogmastigotes and *in vivo* against tripogmastigotes of that tridax shows best antiprotozoal activity.

X. LEISHMANICIDAL ACTIVITY

The Methanol extracts prepared from different parts of 18 plants collected and evaluated in an *in vitro* bioassay for leishmanicidal activity against *Leishmania mexicana* promastigotes. The ten most potent plant extracts (IC₅₀ < 50 µg/ml) were *Aphelandra scabra* leaves, *Byrsonima bucidaefolia* bark, *Byrsonima crassifolia* bark, *Clusia flava* leaves, *Cupania dentata* bark, *Diphyssa carthagenensis* leaves, *Dorstenia contrajerva* whole plant, *Milleria quinqueflora* roots, *Tridax procumbens* whole plant, and *Vitex gaumeri* bark.

XI. ANTI-INFLAMMATORY ACTIVITY

In the carrageenan induced Oedema model, inhibition of Oedema was comparable in 200mg/kg *Tridax procumbens* and 50mg/kg ibuprofen treatment and the specific activity of the enzyme gamma glutamyl transpeptidase was comparable in the *Tridax procumbens*, ibuprofen and aspirin at 200 mg/ kg

XII. CONCLUSION

Although *Tridax procumbens* is type of weed plant it has many useful above discussed properties and activities. The above properties clearly show that it has both immunological and pharmaceutical properties. Some of the beneficial activity are yet to clear about this plant.

XIII. ACKNOWLEDGEMENT

The author expresses their gratitude towards the host Institution Dr.N.G.P. Arts and Science college, Management, Principal, Deans, Head of the department, guide and other all other staffs of Department of biotechnology for rendering all the facilities and support. Communication no: DrNGPASC 2019-20 BS019.

XIV. REFERENCE

- [1] Agrawal SS, Talele GS and Surana S (2009). Antioxidant activity from fractions from *Tridax procumbens*. Journal of Pharmacy Research, 2(1): 71-73
- [2] Ali ME and Ramachandram R (2001). A new flavonoid from the aerial parts of *Tridax procumbens*. *Fitoterapia*, 72(3): 313-315.
- [3] Ali MS and Jahangir M (2002). A Bis-Bithiophene from *Tridax procumbens* L. (Asteraceae). Nat. Prod. Res., 16(4): 217-221.
- [4] Caceres A, Berger I, Barrientos AC, Caceres A, Hernandez M, Rastrelli L, Passreiter CM and Kubelka W (1986). Plants used in Guatemala for the treatment of protozoal infections: II. Activity of extracts and fractions of five Guatemalan plants against *Trypanosoma cruzi*. Journal of Ethnopharmacology, 62, 107-115.
- [5] Devi Vimla and Suneeta A (1990). J. of Research and Education in Indian Medicine, 9(4): 39-41.
- [6] Gadre A and Gabhe SY (1992). Identification of some sterols of *Tridax procumbens* by GC-MS. Indian J. Pharm. Scien., 55(5): 191-192.
- [7] Gupta S, Yadava JNS and Tondon JS (1993). Antisecretory (anti-diarrheal) activity of Indian medicinal plants against *E. coli* enterotoxin- induced secretion in rabbit and guinea pig ileal loop models. International J. of Pharmacology, 31(3): 198-204.
- [8] Kanungo S, Mohanty S, Das M, Patnaik J and Mohanty M (1995). A study of the effects of *Tridax procumbens* Linn. on normal and heparin induced prolongation of clotting time in rabbits. Indian Journal of Pharmacology, 27(1): 63.
- [9] Margaret I, Reddy PS and Kaiser J (1998). Anti-inflammatory profile of *Tridax Procumbens* in animal and fibroblast cell models. Phytotherapy Research, 12(4): 285-287.
- [10] Morales C, Gomez-Serranillos MP, Iglesias I, Villar AM and Caceres A (2001). Preliminary screening of five ethnomedicinal plants of Guatemala. *Farmaco.*, 56(5-7): 523-526.
- [11] Oommachan M (1976). The Flora of Bhopal. I Edn. J. K. Jain Brothers, Bhopal, 192-199.
- [12] Oudhia P (1999). Medicinal weeds in rice fields of Chhattisgarh (India). International Rice Research Note, 24(1): 40-41.
- [13] Pathak AK and Dixit VK (1991). Hepatoprotective activity of *Tridax-procumbens* part I. *Fitoterapia*, 62: 307-314.
- [14] Pathak AK. and Dixit VK (1988). Insecticidal and insect repellent activity of essential oils of *Tridax procumbens* and *Cyathoclineyrate*. *Fitoterapia*, 59: 211-214.



- [15] Peraza-Sanchez SR, Cen-Pacheco F, Noh-Chimal A, May-Pat F, Sima-Polanco P, Dumonteil E, Garcia-Miss MR and Mut-Martin M (2007). Leishmanicidal evaluation of extracts from native plants of the Yucatan peninsula *Fitoterapia*, 78: 315–318.
- [16] Perumal SR, Ignacimuthu S and Patric Raja D (1999). Preliminary screening of ethnomedicinal plants from India. *J. Ethnopharmacol.*, 66(2): 235-240.
- [17] Rajkumar S and Jebanesan A (2007). Repellent activity of selected plant essential oils against the malarial fever mosquito *Anopheles stephensi* Trop. Biomed., 24(2): 71-5.