

# WEED CONTROL IN IMPROVING CROP YIELD WITH TECHNOLOGICAL ADVANCEMENTS

Susanta Kundu  
Faculty of Engineering and Technology,  
Shree Guru Gobind Singh Tricentenary University  
(SGT University), Gurgaon, Haryana, India

Sharif Ahmad  
Former Pro-Vice-Chancellor  
Shree Guru Gobind Singh Tricentenary University  
(SGT University), Gurgaon, Haryana, India

**Abstract:** The need for agricultural production is directly associated with the increased global population and limited agricultural land resources. Natural dependencies on weather & climatic conditions and factors like weeds, diseases, pests, or combinatorial issues play a significant role in the final agricultural yield. While disease and pest controls are relatively straightforward, weed control has more challenges since it coexists with crops as a contender of the same soil nutrition and favorable environment during the plant growth period. Weeds are detrimental to harvest, but it is manageable to protect through various biological, chemical, cultural, or physical mechanisms and compliant with ecological methods. Literature reveals that the weed control methods have few shortcomings, influencing overall crop production. However, the advent of precision technology in agriculture will provide a way to overcome such drawbacks. This article highlights the literature, uses various technological methods to control weeds more precisely and considers other factors influencing weed management. The study also highlights a brief covering the need for management and development of technology involved in weed management, with more thrust on identifying a few more prevailing weed control challenges in India and provision for futuristic planning to support the small farmers as end-users!

**Key Words:** Variable Rate Technology (VRT), Weed Control, weed management.

## I. INTRODUCTION

In the early days, concerning population and availability of abundant agricultural land, the conventional crop growing system satisfactorily met society's needs. However, an exponential increase in population with time and the shrunk-cultivable land area with urbanization led to hunger and

famine as food grain growth per year was poorer than needed. Literature reveals that Indian agricultural development in the era 1950 – 60s was conventional and not backed by any improved processes, causing repeated uncertainty in agricultural growth. However, from the early sixties, with proper government initiatives and the effort of agricultural scientist Professor M. S. Swaminathan had achieved a Green Revolution (GR) by producing high-yielding varieties of rice and wheat to increase food grain production. The Green Revolution tending from the mid-sixties to mid-seventies transformed India from a food-deficient country to a leading agricultural nation [70]. While the initial focus of GR was to expand farming areas, that required awareness and actions in the agriculture field led to the increased crop yields through the use of fertilizers, improved seed quality, chemical pesticides, etc.

Ever since the Green Revolution, India has continued its sustainability in crop production year-on-year rate and higher yield in agriculture and its associated sectors. In 2020-21, it contributed 20.19% Gross Value Product (GVP) (Fig.1) was a measure of whole economic output among three key economic sectors (services, industry, and agriculture)[1].




Fig. 1: GVA share (%) among 3 Keys Economic Sectors in India (2020-21)



The agricultural GDP growth rate in India during 2015-16 to 2019-20[2] is shown in Fig. 2.




Fig. 2: GVA % share of agriculture + related sectors over total Indian Economy

Growth in this sector recorded fluctuating trends over the last few years. There can be many factors responsible for such a trend. One possible reason may be the inadequate soil responses to excessive or inappropriate use of chemical fertilizers [3].

In 2020-21, despite Covid 19, the agriculture sector in India exhibited a positive growth of 20.19% (Fig. 1). Such a variation in agriculture data needs more people's participation and contributions [72].

It is essential to mention that about 54.60% of the total population in India was directly or indirectly associated with agriculture as per the last census in 2011 [2], highlighting the significant involvement of this sector in the Indian socio-economic context. A fall in agricultural land (Fig. 2a) limits crop production due to fast urbanization. It hampers the harvest-growth rate to an alarming extent, coupled with weather & climate as natural dependencies and other factors like weeds, diseases, pests, rodents, or any of these combinations during the cultivation period [71].




Fig 2a: Year-wise Agricultural land (% of land area) - India  
 Source: <https://data.worldbank.org/>

The agricultural yield factor, such as the ratio of crop produced and associated cultivating land, is influenced by weeds as a competition to crop [67], being highly invasive and outcompeting. Tata Strategic Management Group 2014 reported a study [4] on more crop losses due to weed infiltration (34%) when compared with factors like insects

(26%), diseases (26%), rodents (6%), and others (8%) (Fig.3) established a critical role of weeds.




Fig. 3: Criticality of weeds

Information from this study plays a significant role in providing vital information and can be used to generate awareness among the farmers.

The present review article highlights the role of various parameters used for weed control practices improvised by experimental researchers and technological advancements across the globe to achieve higher agricultural yields.

The set policy target of the Government of India to double farmers' income is well in tune with the requirement of the current economic scenario.

The article also emphasizes the impact of technological advances in weed controls. However, these still have a few challenges of acceptance and deployment in developing countries to align as an affordable solution to farmers and cultural acceptance way forward [73].

Weed control practices and research in developing countries Weeds usually grow from the previous season's seed, annual or can regrow perennial from dormant roots, underground stems, tubers, rhizomes, etc. Classification of weeds based on their lifecycle (annual or perennial), origin (indigenous or not), morphology (shoot or root) [74], habitat (terrestrial or aquatic), soil types (black soil, red soil, etc.), or association with a particular crop help to understand weed characteristics for its control, spread and separating from the crop. Weeds can further be identified as invasive weeds that can threaten biodiversity due to their allelopathic effect by producing biochemicals that influence the growth and germination of adjoining plants [75].

Conceptually weed management [34] prevents the growth or spread of weeds through cultural, physical, biological, and chemical methods. Fig. 4 depicts a situational and combinational hierarchy of these weed control methods.




Fig. 4: Available weed control methods

For decades, herbicides have used chemical applications in weed control as a thriving traditional practice [5]. These are applied on the soil surface so that weeds germinating on the top or surface layers get killed due to the herbicide's incidental absorption. For perennial weeds, herbicides are injected as a sub-surface application into the lower soil layers at several points to reach the underground stems, tubers, dormant roots, and rhizomes that kill them [76].

Based on statistical analysis LD50 (Lethal Dose) model of toxicants [77], World Health Organization (WHO) classified insecticides and herbicides as extremely, highly, moderately, and slightly hazardous [6]. Higher toxicity is more detrimental to health and ecological systems [78].

Regulatory controls have been imposed from time to time to meet the sustainability of the environment [79]. Using chemical applications to evolve weed resistance to multiple herbicides opened up studying the genetics of weed adaptation under the biological control of various gene inheritance [7].

Apart from herbicides as chemicals, foliar applications [80] provide fertilizer or chemical nutrients by spraying selectively on crop leaves, thereby depriving weeds of such nutrition [81].

Biological methods [8] deploy specific insects, pathogens, or animals acting as bio agents to invade weeds and prevent their growth or kill them. Traditionally, biological controls play a significant role in reducing weeds but not eradicating them. Physical methods apply physical forces to pull out and kill weeds through the human, animal, or mechanical tools. Such as hand-weeding, hand-hoeing, digging, burning, etc., are widely practiced depending on weed and crop situation during a cultivation period. Cultural practices help in creating favorable crop conditions. For example, regular cleaning and maintaining the sanity of crop area through tillage, irrigation, adding fertilizer or nutrients closer to plants, crop rotation, etc., are some of the popular cultural practices used in weed control. In addition, the

removal of aquatic weeds is a preventive process applied to avoid water loss and insufficient water flows. It improves the physical & chemical characteristics of both water and hydro soil [9].

After sowing seeds, weeds start growing and competing with crops simultaneously. Thus Integrated Weed Management (IWM) is a situational approach [82] used to manage weeds infields by incorporating techniques or available methods as and when needed. It best uses the prevailing situation in a crop growing season. Chemical & biological prevention processes [83] are widely practiced in India through labor-intensive care. The use of technical gadgets can be an alternative to some of these methods.

Researchers, for better control of weeds and reduction in the use of herbicides/chemicals, experimented with different soil qualities. It is also observed that the physical weed control mechanisms such as precise inter and intra-row cultivation can reduce the use of chemicals.

Central Research Station, College of Agriculture in Bhubaneswar studied the performance and role of nutrients and microbial contents of soil on crop growth at the end of two consecutive seasons using combinations of different weed management and varying herbicide/chemical concentration. It improved soil quality through a microbial population of bacteria, fungus, actinomycetes, and dehydrogenase activities [8].

Anand Agricultural University (Gujarat) experimented on irrigated wheat using combinations of herbicide treatments during two successive rabi seasons. It yielded the required weed density and superior wheat grains due to the post-emergence application of combinatorial herbicides [10]. Further, the analysis of two years of rice production data at the Agricultural Research Station, Ragolu (Andhra Pradesh), revealed a combination of green manure and need-based hand weeding yielded an increased benefit to cost ratio [11].

Tabuk University – Saudi Arabia conducted a study on the role of allelopathic compounds using natural herbicides on seed germination that resulted in higher weed control [12]. Such natural and eco-friendly approaches can supplement improved weed control by finding compounds that inhibit crop seeds and prevent unwanted weed germination.

Agronomy researchers at Kumarganj, Ayodhya, used a combination of nitrogenous compounds as weed control treatments on turmeric. In two successive years, they recorded improved plant attributes like plant height, higher leaf area index (LAI), lower value of weed density, and dry mass [13].

Zonal Agriculture Research Station, Kalaburagi (Karnataka), studied the influence of weed management by allelopathy practices on nutrient status using plant extracts with a 50% reduction in herbicide. It suggests an increase in soil fertility and productivity over an extended period. The resultant outcome also indicates a more productive and sustainable growth [14].



The experimental results on using pre-emergent herbicides suggested the killing of weeds before sprouting. It could act as an effective weed control method among all possible options, such as no weed control, manual weeding, and pre-emergent and post-emergent herbicide treatments [15]. A similar study on an irrigated blackgram at the College of Agricultural Technology, Kullapuram (Tamil Nadu), revealed the effects of pre-emergence (PE) and early post-emergence (EPoE) use of herbicides on weed management [16].

One of the studies experimented on a drill-cum-weedicide applicator developed in an indigenous tractor-operated seed sowing machine. Improvised mechanical tools used in herbicide applications reduced spraying time, material waste, and chemical exposure to farmers [17].

Understanding the dynamism in weed biology [84] is one of the important factors playing a significant role in developing and improving techniques, tools, and processes in managing agricultural yields. Furthermore, data and research results are essential to rolling back to the field for appropriate weed control mapping to climate (temperature, humidity, etc.) [85] and microbial soil conditions [40].

**Use of herbicides in weed control**

Herbicides are an essential part of agriculture due to their higher efficiency in weed control [39], improving crop production, and cost-effectiveness. Herbicides can be both inorganic and organic. Inorganic herbicides are synthetic, created in labs, while organic ones are made from natural chemicals. Organic herbicides have low toxicity, break down quickly and leave minimum residual effects, providing environmental benefits by being biocompatible [86]. High-potency herbicides that work in low doses are in demand to control broad-spectrum weeds [4].

Widespread use of herbicides exploited its efficiency in weed control resulting in soil quality deterioration [87] and leaving environmental impacts due to residual effects.

Recent progress of nanotechnology in the agricultural domain is ushering a new hope in herbicide usage through nano-herbicides for a sustainable and eco- friendly solution. It will supplement the end-users to control weeds using much lesser toxicity and chemicals, increasing manifold benefits [18].

Nearly 270 herbicides were added globally between 1936 and 2012 [4], as shown in Fig. 5.




Fig. 5: Herbicide counts vs. Year of Introduction  
 Source: Pesticide Properties DB, University of Hertfordshire, 2015

Not many new chemicals were added post-nineties, indicating an overall downward trend with time. India imported herbicides way back in the 1960s, primarily for tea plantations [88].

However, herbicides drew the attention of farmers with the increase in labor costs for manual weed removal. Farmers switched to herbicides for higher efficiency and improved cost-effectiveness.

The consumption soon increased manifold. Nearly 700 formulations of herbicides are available in the Indian market to cater to weed control [4].

In agriculture, insecticides, herbicides, bactericides, fungicides, and rodenticides are collectively called pesticides. Thus, the herbicides are a subset of pesticides.




Fig.6: Herbicide usage pattern in India during 1990 - 2010

The use of aggregated pesticides and herbicides in India shows (Fig. 6 and 7) a downtrend in consumption till 2008. Increased and ever-rising labor costs and labor scarcity added by technological intrusions accelerated higher consumption of herbicides.



Fig. 7: Aggregated (1990 - 2010) Pesticide (total) use trend in India




Fig. 8: Aggregated (1990 - 2010) top 10 herbicide Aggregated herbicide usages from 1990 to 2010, shown in Fig. 8, have relative positions in the top 10 countries.

It is reflected in the increased usage pattern in post-2008, substantiating projected herbicide consumption in India growing at a CAGR of 15% as planned till 2020[19].

Table 1 (Source: FAOSTAT) captures the herbicides used in India from 1990 to 2010.

Table 1: Herbicides used in tones during 1990 - 2010 in India								
Year	Amides	Bipiridils	Dinitroanilines	Phenoxy hormone products	Triazines	Urea derivates	Other	Total
1990				791	275	1804	2951	5821
1991				686	285	1881	3151	6003
1992				750	277	2290	4120	7437
1993	1113			629	310	2307	2234	6593
1994	1919	46	95	612	472	2568	1098	6810
1995	2271	25	94	467	207	2490	566	6120
1996	2596	35	114	621	353	2657	882	7258
1997	2644	41	112	609	350	2674	562	6992
1998	2647	62	164	659	452	2662	835	7481
1999	2598	113	161	680	424	2685	819	7480
2003			213			2842	4445	7500
2004			115			2333	2706	5154
2005			289			1447	5223	6959
2006			82			2251	3971	6304
2007			98			1428	2596	4122
2008			81			1217	2276	3574
2009			49			1215	2956	4220
2010			71			1834	4430	6335
Total	15788	322	1738	6504	3405	38585	45821	112163

Table 2 (Source: FAOSTAT) has comparative data on top herbicides and pesticide-consuming countries with indicative percent use of herbicide/pesticides.

Country	Pesticides Use (tones)	Herbicides (tones)	(Herbicides / Pesticides) %
USA	407,052	202,493	49.7%
Brazil	166,048	94,868	57.1%
USSR	89,200	40,100	45.0%
France	87,806	29,651	33.8%
Colombia	54,770	25,205	46.0%
India	47,698	6,411	13.4%



Figures 9 and 10 show India's harvested land and crop production from 1961 to 2019 for rice and wheat, respectively (Source: FAOSTAT). The growths are aligned with the onset of the green revolution, using fertilizers, pesticides, and herbicides to improve production yield. Usual monsoon rain, good soil moisture, and support from the government encourage farmers to favor growing rice and wheat as the main crop in India [20]. The data indicates near-linear growth with harvest area in Fig 9, 10. However, in recent years – particularly post-2015- rice and wheat production rates rose even higher despite a partial dip in harvest areas – primarily due to urbanization.




Fig. 9: Production / Yield quantities of Rice in India during 1961 to 2019

Figs 9 and 10 also substantiate consistent growth and high crop yields from the green revolution period for rice and wheat, respectively. Constant effort to educate farmers on the correct use of fertilizers, herbicides, application frequency, dosages, etc., resulted in aligning such growth rates [89]. However, reaching the farmers for such instances is not so easy. It is also challenging for the farmers to convey their needs effectively [19].

### Technological advancements

Most small farmers usually pluck weeds manually from the field or spray herbicides using manual backpacks. Their weed removal tools include brush cutter, trimmer, power weeders, etc. Backpacks prayers are vulnerable due to the lack of precision in the quantity needed and health concerns of persons involved in spraying. Existing practices have low yields without appropriate technology for precision usage. With an intent to increase the farmers' earnings, smart agriculture with intelligent technologies is essential to bring a difference. IoT-based innovative applications added a revolution to the traditional weed control methods, supported by GPS-based remote-controlled robots with smart onboard sensors, Wi-Fi, and cameras [21]. Data from soil sensors, temperature & humidity sensors, air quality sensors, and sensors capturing video and

photographs can be monitored near real-time and processed for decision making [90].




Fig. 10: Production / Yield quantities of Wheat in India during 1961 to 2019

Robotics in weed management is supported by computer vision and AI-triggered robotic actions precision in weed control using a programmable expert system to control and manage its activities [38] automatically. Variable Rate Technology (VRT) enables an automated decision-making robot to apply herbicides at variable rates without manually changing equipment. VRT enables Variable Rate Applications (VRA) of materials to a given landscape based on sensors, maps, and GPS data. The system can ensure environmental safety [22] by controlled use of herbicides.

Combined with electronics and mechanical engineering – mechatronic scan work as an autonomous machine for weed control in the field [43]. Statistical tools and analysis of captured images by guided drones can help balance herbicide spray and weed removal by a trained robot with sensors mounted on it [23]. Herbicides control the weeding problem and increase crop growth [37].

Table 3: Source <https://datacommons.org>

Rank	Country	Population (2019)
1	People's Republic of China	1,396,530,000
2	India	1,254,020,000
3	United States of America	328,016,242
4	Indonesia	267,871,000
5	Pakistan	216,565,318
6	Brazil	208,495,000
7	Nigeria	200,963,599
8	Bangladesh	163,046,161
9	Russia	146,675,000
10	Japan	126,443,000

Robotics in weed control to hit precision targets need spatial localization [36] of weeds for precision application of herbicide spray in Site-Specific Weed Management (SSWM) using Unmanned Aerial Vehicles (UAV). It can monitor weeds and provide decision-supporting information



for precision spraying. Results demonstrated increased efficiency with competitive accuracy [24].

The development of proximal imaging systems with sensors closer to objects, when embedded in autonomous vehicles, can estimate plant morphology better above the ground dry matter biomass (BM) and leaf area index (LAI) at early growth stages [25]. Satellite-based remote sensing detects and maps weed presence through broadband multispectral sensors like VNIR (Visible and Near Infrared) to derive Atmospheric Resistance Vegetation Index (ARVI) as fundamental variables [26].

Improvement in weed management over traditional methods, transformed with the advancement of IoT use of remote-controlled robotic vehicles, low power, and low-cost wireless sensors. Related notifications on smart phone applications engage farmers using Wi-Fi, 3G, 4G/LTE [27]. Mobile phones are now an essential communication tool that empowers farmers with appropriate information in weed management [28]. Institutional roles are necessary for knowledge dissemination through such devices.

Table 4: Source <https://datacommons.org>

Country	Population Growth Rate (%) 2019
India	1.02%
United Kingdom	0.56%
Germany	0.27%
Canada	1.42%
People's Republic of China	0.36%
United States of America	0.47%

**Challenges**

India is estimated to have nearly 18% of the world's population [29]. In 2019, the population in India ranked second in the world next to China (Table 3). An increasing population (Table 4) and ever-increasing urbanizations require increased agricultural production as the focus of all farmers.

Weed control is an optimizing operational factor to enhance the yield. The challenge stretched further to emphasize food grain self-sufficiency and improve food security [33].

Applications of herbicides have inherent ecological challenges. Farmers at the grass-root level should be aware of such acts. Results of intrusive research in agriculture should be spread to farmers as best practices to cover the issues under discussion.

Combinatorial use of hybrid herbicides deployed using precision tools is some improvisations or extended practices to reduce the impacts of weed control on ecology. However, such technical fixes or improvements are yet to align and address the vulnerability of herbicide-resistant weeds, environmental impacts, and public health concerns.

Dynamism in the simultaneous growth of weeds and crops continues to be a critical factor in agricultural yield loss. Scientific understanding of weed biology and related competition with crops for an integrated implementation approach may help higher yield resolution.

Many farmers use backpacks to spray herbicides for weed control [42]. Such manual processes involve handhold sprayers and a manually operated compressed cylinder of liquid chemicals. This process induces uniform application of chemicals on visible weed areas with no scope to apply to specific coverage regions. A precision spraying-weeders for optimal herbicide use can reduce herbicide consumption by 80% and reduce cost. Such improvement will minimize environmental risk and supplement economic farming. It's important to share such information with the farmers as best agricultural practices and demonstrate improved mechanisms to achieve it.

For more extensive cultivating lands, robots in agriculture and robotic technology ease weed management. However, it has challenges related to crop types, the shape of the field, plant spacing, etc., that should consider before evaluating such options. For example, a robotic lawnmower (RLM) has a better working performance when compared with a riding mower (RM) and a walking mower (WM) in an orchard [30].

The challenge is to deploy efficient methodologies catering to site-specific weed control.

Precision tools with robotics and image processing in weed control are not available with easy affordability for sustainable solutions in many places, including India. Most farmers with smaller land areas depend on available cost-effective tools with low efficient conventional processes. Moreover, these tools do not address ecological and related harms, though they try to minimize them.

Industry involvement in device-advanced sensor technology in precision weed control at an affordable cost is a challenge. There will be hurdles in farmer awareness of such advanced devices' safe and convenient use. Thus, training is needed to use technology appropriately to improve crop growth rate. Predicting crop yield with weed factor will continue to be tricky, though wider acceptance and deployment of technical development can ease it further.

**The way forward for futuristic scope**

The use of modern technology in weed control helped save the crops and improve the overall production rate. The role of advanced technology in increasing production rate is recorded or observed. It reflects that using advanced instruments or devices for herbicide applications in agriculture may provide a good future scope in enhancing



crop production rates. Combinatorial use of herbicides using precision tools improves existing practices to reduce the impacts of weed control on ecology (resistance, ecology, health, etc.). Systems-level ecological thinking can help appropriate agronomical decisions to achieve sustainable weed management.

However, due to the steep increase in population, the agricultural production rate may not always match society's needs. As discussed, many risks and challenges are still associated with weed control. To overcome such challenges, researchers are working in multiple directions. Weed characteristics differentiating from crops are intensifying. Researchers can address the impacts of herbicide residues, herbicide-resistant weeds, the results of global climate change on crop-weed interaction, and environmental safety. With awareness of higher efficiency in weed control, herbicide consumption is rising. Dissemination of information to end-users to create awareness of ecological pressure is crucial. Impacts demonstrated to minimize herbicidal residues by local organizations through academic and research institutions collaborating with local administrations (panchayat, etc.) will directly connect to end-users (the farmers).

To balance technological advances and catering to the benefits through appropriate tool deployment among ordinary farmers. It's time to support entrepreneurial efforts in developing various applications and technical instruments for weed control using advanced wireless sensors targeted at small farmers and providing services to them. The approach may open a vista for new agricultural entrepreneurship in rural development.

These discussions suggest that a proper application of weed control technology can improve the required growth rate of agricultural production over time which will significantly enhance the level of society.

## II. CONCLUSION

The present review article aims to discuss the literature covering various aspects of applications in technology in controlling weeds and enhancing crop growth. The article also covers various technological improvisations and applications in weed control.

Literature reports the harmful impacts of weeds on agricultural growth. It also highlighted the role of technology in overcoming the various challenges caused by weeds in the agro-economy and crop production. In view of this, the proposed review article discusses crop production and its salient features, constrained by natural factors (weather & climate) and factors like weeds, diseases, pests, rodents, etc.

The thrust of the review is to highlight the various parameters that influence the crop production cultivating

land ratio. The agricultural yield factor measures the percentage of crop produced and associated enabling land area. While weather and climate are beyond human control, there is an utmost need to control weeds through improvisations in weed control processes and related applicable devices. A better grip over the issues requires a good understanding of weed characteristics in controlling the spread and separating it from the desired crop through appropriate processes. It can prevent weeds from growing or spreading through cultural, physical, biological, and chemical methods.

For decades, the chemical method has been the most successfully practiced using various herbicides among all the weed control practices. Such weed control approaches have a few shortcomings that need technological improvements to influence crop production with maximum green and salty land utilization as an agricultural field. Developments in precision technology and related alignment with agricultural applications aimed at overcoming existing drawbacks and challenges of prevailing weed control methods.

Agricultural growth impacted by weeds, associated challenges, and futuristic scopes highlighted in this review suggests that technology plays a significant role in the agricultural growth rate to meet society's food security. It also allows a country to become self-sufficient in crop growth with available agricultural land.

## III. REFERENCES

- [1]. Sector-wise G.D.P. of India, Source: Ministry of Statistics and Program Implementation, Date: 17 June 2021  
<https://statisticstimes.com/economy/country/india-gdp-sectorwise.php>
- [2]. Agricultural AR 2020-21 (Hindi) dated 06 April 2021 pg  
[1https://agricoop.nic.in/sites/default/files/approved%20AR-Hindi%20%281%29.pdf](https://agricoop.nic.in/sites/default/files/approved%20AR-Hindi%20%281%29.pdf)
- [3]. Soil pollution caused by inappropriate use of chemical fertilizers Maria-Alexandra Enescu. Vrije Universiteit Brussel November 2019 DOI:10.13140/RG.2.2.35325.51688 [https://www.researchgate.net/publication/340032701\\_SOIL\\_POLLUTION\\_CAUSED\\_BY\\_INAPPROPRIATE\\_USE\\_OF\\_CHEMICAL\\_FERTILIZERS](https://www.researchgate.net/publication/340032701_SOIL_POLLUTION_CAUSED_BY_INAPPROPRIATE_USE_OF_CHEMICAL_FERTILIZERS)
- [4]. Herbicide Use in Indian Agriculture – ICAR, Information Bulletin No. 22 by Partha P. Choudhury, Raghendra Singh, Dibakar Ghosh, A.R. Sharma <https://krishi.icar.gov.in/jspui/bitstream/123456789/614/1/Information%20Bulletin%20No%20-%202022%20->





- [%20Herbicide%20Use%20in%20Indian%20Agric ult ure.pdf](#)
- [5]. WHO Recommended Classification of Pesticides 2004 [https://www.who.int/ipcs/publications/pesticides\\_hazard\\_rev\\_3.pdf](https://www.who.int/ipcs/publications/pesticides_hazard_rev_3.pdf)
- [6]. The Role of Population and Quantitative Genetics and Modern Sequencing Technologies to Understand Evolved Herbicide Resistance and Weed Fitness Ramon G Leon; Jeffrey C Dunne; Fred Gould - Journal List Wiley-Blackwell Online OpenPMC7754128 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7754128/>
- [7]. Soil Biological and Biochemical Properties in Rice-Maize System under Different Nutrient and Weed Management Practices by BanashriLodh;S N Jena;R K Paikaray <https://www.ijcmas.com/10-1-2021/Banashri%20Lodh,%20et%20al.pdf>
- [8]. Environmental Assessment of Aquatic Weeds, Water and Sediments Related to Rehabilitation and Manual Maintenance (Case Study: Desonas Canal, Egypt) Yasser Mahmoud Ali [http://www.lifesciencesite.com/lsj/lsj170420/05\\_362\\_59lsj170420\\_22\\_31.pdf](http://www.lifesciencesite.com/lsj/lsj170420/05_362_59lsj170420_22_31.pdf)
- [9]. Management of Complex Weed Flora through Herbicide Combinations in Irrigated Wheat (*Triticum Aestivum* L.). Vipul Y Patel;B D Patel;V J Patel;M B Viradiya;H B Sodavadiya <https://www.ijcmas.com/10-1-2021/Vipul%20Y.%20Patel.%20et%20al.pdf>
- [10]. Organic Weed Management in Transplanted Rice K V Ramana Murthy;A Upendra Rao;D Adilakshmi <https://bioinfopublication.org/includownload.php?id=BIA0005610>
- [11]. Herbicidal Effects of Prosopis Juliflora (Sw.) D.C. (Mesquite) Leaf Powder on Seed Germination and Seedling Growth of the Weed Species *Tribulus Terrestris* L Abdelrahim Adam Elbalola <http://www.sciencewebpublishing.net/jacr/archive/2020/September/pdf/Elbalola.pdf>
- [12]. Effect of Nitrogen Sources and Weed Management Practices on Weed Dynamics and Yield of Turmeric Prashant Singh; Ravi Shanker Singh; Dinesh Kumar; Rajesh Kumar; Sanjay Kumar; Mahendra Pratap Singh; Gajendra Singh <https://www.phytojournal.com/archives/2020/vol9issue5/PartI/9-5-87-309.pdf>
- [13]. Influence of Weed Management by Allelopathy on Growth, Yield and Nutrient Status of Soil in Pigeonpea (*Cajanus Cajan* L.) Shridhara; B Nagoli; B M Dodamani; M Y Ajaya Kumar; Pandit S Rathod; Mahadeva Swamy; <https://www.ijcmas.com/10-1-2021/Shridhara,%20et%20al.pdf>
- [14]. Assessment of Different Weed Control Methods on Growth and Yield Performance of T. Aus Rice Muhammad Asif Osman, KawsarHossen, RabiulHaq Chowdhury, Chowdhury Nafisa Tabassum and Md Khairul Islam [https://juniperpublishers.com/artoaj/ARTOAJ\\_MS.ID.556267.php](https://juniperpublishers.com/artoaj/ARTOAJ_MS.ID.556267.php)
- [15]. Effect of Pre and Early Post Emergence Herbicides on Weed Control in Irrigated Blackgram (*Vigna mungo* L.) by Sukumar J; Kunjammal P <http://masujournal.org/107/190344.pdf>
- [16]. Development of Tractor Operated Seed Ferti Drill-Cum-Weedicide Applicator A S Ghadge; A S Waghmode; K B Phalphe; S H Thakare; A K Kamble <https://www.ijcmas.com/9-2-2020/A.%20S.%20Ghadge.%20et%20al.pdf>
- [17]. Nanotechnology in Herbicide Resistance - Evy Alice Abigail and Ramalingam Chidambaram - <https://www.intechopen.com/chapters/55009>
- [18]. National Round Table of C.E.O.s on Indian Plant Protection - Indian Council of Food and Agriculture 17 Feb 2017 [https://www.icfa.org.in/assets/doc/reports/sou\\_pp.pdf](https://www.icfa.org.in/assets/doc/reports/sou_pp.pdf)
- [19]. Grain and Feed Annual – 2021 USDA - Report Number: IN2021-0040, 31 Mar 21 / <https://apps.fas.usda.gov/newgainapi/api/Report/DownloadReportByFileName?fileName=Grain%20and%20Feed%20Annual%20New%20Delhi%20India%2003-15-2021.pdf>
- [20]. Internet of Things (IoT) Based Polyhouse Monitoring and Controlling by Gholap Rohit Ramesh; Malwade Akash Gorakshnath; Shinde Nitin Pradeep; Warale Shradha <http://www.ijrsd.com/articles/IJSRDV7I20777.pdf>
- [21]. Challenges of Multi/hyper Spectral Images in Precision Agriculture Applications – by Adriano Mancini; Emanuele Frontoni And Primo Zingaretti <https://iopscience.iop.org/article/10.1088/1755-1315/275/1/012001>
- [22]. Drone and Sensor Technology for Sustainable Weed Management: A Review – by Esposito, Marco; Crimaldi, Mariano; Cirillo <https://chembioagro.springeropen.com/articles/10.1186/s40538-021-00217-8>
- [23]. Lightweight Semantic Segmentation Network for Real-Time Weed Mapping Using Unmanned Aerial Vehicles Jizhong Deng; Zhaoji Zhong; Huasheng Huang; Yubin Lan; Yuxing Han; Yali Zhang <https://www.mdpi.com/2076-3417/10/20/7132>
- [24]. Rgb Image-Derived Indicators for Spatial Assessment of the Impact of Broadleaf Weeds on Wheat Biomass Christelle Gée; Emmanuel Denimal <https://www.mdpi.com/2072-4292/12/18/2982>



- [25]. A Comparative Analysis of Planetscope and Sentinel-2 Space-Borne Sensors in Mapping Striga Weed Using Guided Regularised Random Forest Classification Ensemble B T Mudereri; T Dube; E M Adel-Rahman; S Niassy; E Kimathi; Z Khan; T Landmann <https://www.int-arch-photogramm-remote-sens-spatial-inf-sci.net/XLII-2-W13/701/2019/isprs-archives-XLII-2-W13-701-2019.pdf>
- [26]. IoT in Agriculture by Nanda M B; Sharath <http://www.ijsrd.com/articles/IJSRDV7I30693.pdf>
- [27]. Seeking of Agriculture Information through Mobile Phone among Paddy Farmers in Selangor Nor Sabila Ramli; Md Salleh Hassan; Norsida Man; Bahaman Abu Samah; Siti Zobedah <https://hrmars.com/papers/detail/IJARBS/5969/Seeking-of-Agriculture-Information-through-Mobile-Phone-among-Paddy-Farmers-in-Selangor>
- [28]. India – Population <https://www.globalsecurity.org/military/world/india/population.htm>
- [29]. Weed Management and Economic Analysis of a Robotic Lawnmower: A Case Study in a Japanese Pear Orchard, by Muhammad Hossain; Masakazu Komatsuzaki <https://www.mdpi.com/2077-0472/11/2/113>
- [30]. Invasive Johnsongrass, a Threat to Native Grasslands and Agriculture by Klein, Peter; Smith, Charles Michael <https://link.springer.com/article/10.2478/s11756-020-00625-5>
- [31]. Effect of Different Scented Rice Variety, Planting Geometry and Weed Management Practices on Yield Parameters and Yield at Southern Bastar of Chhattisgarh, by Purnima Haldar; Ashwani Kumar Thakur; Manish Kumar; G P Ayam; D P Singh <https://www.ijcmas.com/10-1-2021/Purnima%20Haldar,%20et%20al.pdf>
- [32]. Projection for Plant Protection Machinery of Kayseri Province by Necati Cetin; Bunyamin Demir; Cevdet Saglam <https://dergipark.org.tr/en/download/article-file/1116829>
- [33]. Worldwide pesticide usage and its impacts on ecosystem - by Anket Sharma, Vinod Kumar, Babar Shahzad, Mohsin Tanveer, Gagan Preet Singh Sidhu, Neha Handa, Sukhmeen Kaur Kohli, Poonam Yadav, Aditi Shreeya Bali, Ripu Daman Parihar, Owias Iqbal Dar, Kirpal Singh, Shivam Jasrotia, Palak Bakshi, M. Ramakrishnan, Sandeep Kumar, Renu Bhardwaj & Ashwani Kumar Thukral - SN Applied Sciences volume 1, Article number: 1446 (2019) <https://link.springer.com/article/10.1007/s42452-019-1485-1>
- [34]. Influence of Weed Management Practices on Weed Control Efficiency, Yield and Economics in Sweet Corn (*Zea Mays L. Saccharata*) - Bhardwaj S SS; Mishra G C; Lenka S K, <http://www.indianjournals.com/ijor.aspx?target=ijor:ijaeb&volume=13&issue=1&article=005>
- [35]. Allelopathic Effect of Siam Weed Debris on Seed Germination and Seedling Growth of Three Test Crop Species Mahfuza Begum; Abdus Salam; Farhana Zaman <https://actascientificamalaysia.com/archives/ASM/2asm2020/2asm2020-27-32.pdf>
- [36]. Spatial Distribution of Weed Seed Banks in the Agricultural Field and Anthropized Cerrado Klayton Antonio Do Lago Lopes; Marcelo De Sousa Da Silva; Leandro Dos Santos Costa; Taciella F <http://www.macrothink.org/journal/index.php/jas/article/view/16031/12992>
- [37]. Growth Analysis of some Maize Cultivars on Weedy Condition by M Rahayu; PYudono; DIndradewa; EHanudin <https://iopscience.iop.org/article/10.1088/1755-1315/653/1/012075>
- [38]. The use of agricultural robots in weed management and control by Brian L. Steward, Jingyao Gai, Lie Tang [https://lib.dr.iastate.edu/cgi/viewcontent.cgi?article=2333&context=abe\\_eng\\_pubs](https://lib.dr.iastate.edu/cgi/viewcontent.cgi?article=2333&context=abe_eng_pubs)
- [39]. Weed control efficiency of herbicide sulfosulfuron in potato crop – by SP Singh, S Rawal, VK Dua, and SK Sharma1 - Potato J 44 (2): July - December, 2017 [https://www.researchgate.net/publication/323167217\\_WEED\\_CONTROL\\_EFFICIENCY\\_OF\\_HERBICIDE\\_SULFOSULFURON\\_IN\\_POTATO\\_CROP](https://www.researchgate.net/publication/323167217_WEED_CONTROL_EFFICIENCY_OF_HERBICIDE_SULFOSULFURON_IN_POTATO_CROP)
- [40]. Studies about the Soil Microbial Populations, Root Development and Economics of Paddy as Influenced by Sea Weed Extract (*Ascophyllum nodosum*) Rajeev Kumar; Ram Pyare; Uday Pratap Singh; Yatindra Kumar; Anjani Kumar Singh <https://www.phytojournal.com/archives/2020/vol9issue5/PartI/9-5-87-309.pdf>
- [41]. Comparative Assessment of Multiple Linear Regression and Fuzzy Linear Regression Models by Pandit, Pramit; Dey, Prithwiraj; Krishnamurthy, K N <https://link.springer.com/article/10.1007/s42979-021-00473-3>
- [42]. A Review on Agriculture Spraying with Weeding Process Keval Panera R; Jaydeep Jogel N; Pratik Chhaniyara N; Jigar Bhalodiya V; Jagdish Bhakhar D; Bhavik [http://www.ijirset.com/upload/2021/february/30\\_A\\_REVIEW.PDF](http://www.ijirset.com/upload/2021/february/30_A_REVIEW.PDF)
- [43]. Heat-Applicator Machine for Weed Control Spagnolo, Roger T; Custódio, Tiago V; Morais, César S De; Reis, Ângelo V Dos; Machado, Antônio L <https://www.scielo.br/pdf/eagri/v40n5/1809-4430-eagri-40-05-0595.pdf>
- [44]. Agricultural Weeds Extinguisher with the Use of ML Hemanath; Hariharan;



- Aravindhan <https://www.irjet.net/archives/V7/i10/IRJET-V7I10174.pdf>
- [45]. A Review on Inter-Row Crops Mechanical Weeder – by Adewale Sedara; Oluwadunsin Sedara <https://dergipark.org.tr/en/download/article-file/1168827>
- [46]. Evolution of Smart Strategies and Machines Used for Conservative Management of Herbaceous and Horticultural Crops in the Mediterranean Basin: A Review – by Sabina Failla; Michel Pirchio; Mino Sportelli; Christian Frasconi; Marco Fontanelli; Michele Raffaelli; Andrea Peruzzi <https://www.mdpi.com/2073-4395/11/1/106>
- [47]. Dataset of Annotated Food Crops and Weed Images for Robotic Computer Vision Control Kaspars Sudars; Janis Jasko; Ivars Namatevs; Liva Ozola; Niks Badaukis <https://www.sciencedirect.com/science/article/pii/S2352340920307277?via%3Dihub>
- [48]. Determinants of Smallholder Maize Farmer's Perception on Use of Improved Weed Control Technologies in Eswatini Khumalo M; DKibirige; M B Masuku; H R Mloza-Banda; W O Mukabwe; B P Dlamini <http://www.ccsenet.org/journal/index.php/jsd/article/view/0/42753>
- [49]. Experimental Efficiency Evaluation of 445 Nm Semiconductor Laser for Robotized Weed Control Applications by Osadcuks V; Kostromins A; Pecka A; Kotelenecs V; Jasko J [https://agronomy.emu.ee/wpcontent/uploads/2020/05/AR2020\\_Vol18SI2\\_Osadcuks.pdf#abstract-7639](https://agronomy.emu.ee/wpcontent/uploads/2020/05/AR2020_Vol18SI2_Osadcuks.pdf#abstract-7639)
- [50]. Sensor Based Automatic Irrigation for Computer Crops Water Requirements Using Robotic Technology by Kalyanasundaram P; Anto Bennet M; Giridharreddy K <http://sersc.org/journals/index.php/IJCA/article/view/16370/8227>
- [51]. Roboculture Web Based Multipurpose Agricultural Robot by Neena Elias Bardeskar; Nikita Mahadev Surve; Rajlaxmi Ajit Raorane; Ashish Ganotra <http://www.ijrsd.com/articles/IJSRDV7I110422.pdf>
- [52]. Applications of Mechatronics and Autonomous Systems in Smart Agriculture: Potentials and Challenges for Nigeria by Ozoemena Ani; Kennedy Chukwuma; Gideon Okoro; Anselm Oyiga [http://www.aeuso.org/includes/files/articles/Vo110\\_Iss35\\_4502-4517\\_Applications\\_of\\_Mechatronics\\_and\\_Au.pdf](http://www.aeuso.org/includes/files/articles/Vo110_Iss35_4502-4517_Applications_of_Mechatronics_and_Au.pdf)
- [53]. Advanced Machine Learning in Point Spectroscopy, Rgb-And Hyperspectral-Imaging for Automatic Discriminations of Crops and Weeds: A Review Wen-Hao Su <https://www.mdpi.com/2624-6511/3/3/39>
- [54]. Iot Based Energy Efficient Routing Strategy in Wireless Sensor Networks for Precision Agriculture by Muruganandam K; Usha Chauhan [http://www.irdindia.in/journal\\_ijraet/pdf/vol8\\_iss8/10.pdf](http://www.irdindia.in/journal_ijraet/pdf/vol8_iss8/10.pdf)
- [55]. Agricultural Robots: The Applications of Robotics in Smart Agriculture: Towards more Advanced Agro Informatics Practice – by Paul P K; Sinha R R; Aithal P S; Ricardo Saavedra; Bashiru Aremu; Mewada S <https://www.trp.org.in/wp-content/uploads/2020/11/ARME-Vol.9-No.1-January-June-2020-pp.38-44.pdf>
- [56]. Convergence of Wireless Sensor Network and Data Mining for Pest Management in Agriculture by Deepa B; Nandhini S; Jeen Marseline K S <https://www.ijrte.org/wp-content/uploads/papers/v7i6s5/F10960476S519.pdf>
- [57]. Artificial Intelligence-Driven Autonomous Robot for Precision Agriculture by Beloev, Ivan; Kinaneva, Diyana; Georgiev, Georgi; Hristov, Georgi; Zahariev, Plamen <https://www.sciendo.com/article/10.2478/ata-2021-0008>
- [58]. Timing of weed management and yield losses due to weeds in irrigated rice in the Sahel by D.E. Johnson, M.C.S. Wopereis, D. Mbodj, S. Diallo, S. Powers, S.M. Haefele [https://www.researchgate.net/publication/223870524\\_Timing\\_of\\_weed\\_management\\_and\\_yield\\_losses\\_due\\_to\\_weeds\\_in\\_irrigated\\_rice\\_in\\_the\\_Sahel](https://www.researchgate.net/publication/223870524_Timing_of_weed_management_and_yield_losses_due_to_weeds_in_irrigated_rice_in_the_Sahel)
- [59]. Selecting Appropriate Weed Control Systems for Developing Countries by Douglas Young, Stanley Miller, Herbert Fisher and Myron Shenk <https://www.jstor.org/stable/4042700>
- [60]. Weeds as a medical plant: Cynodon dactylon <https://scialert.net/fulltext/?doi=rjnp.2011.508.514>
- [61]. Weeds as a medical plant: Cyperus rotundus <https://curejoy.com/content/health-benefits-musta-cyperus-rotundus/>
- [62]. WSSA POSITION STATEMENT ON BIOLOGICAL CONTROL OF WEED <https://wssa.net/wssa/weed/biological-control/>
- [63]. An ecological future for weed science to sustain crop production and the environment. A review Chloe MacLaren, Jonathan Storkey, Alexander Menegat, Helen Metcalfe, Katharina Dehnen-Schmutz <https://link.springer.com/content/pdf/10.1007/s13593-020-00631-6.pdf>
- [64]. Vision 2050 - Indian Council of Agricultural Research (ICAR) <https://icar.org.in/files/Vision-2050-ICAR.pdf>
- [65]. Economic Survey 2018-19: Report Summary <https://www.prsindia.org/report-summaries/economic-survey-2018-19>



- [66]. India Budget Vol. 2  
[https://www.indiabudget.gov.in/economicsurvey/doc/vol2chapter/echap07\\_vol2.pdf](https://www.indiabudget.gov.in/economicsurvey/doc/vol2chapter/echap07_vol2.pdf)
- [67]. Grand Challenges in Weed Management by Bhagirath Singh Chauhan  
<https://www.frontiersin.org/articles/10.3389/fagro.2019.00003/full>
- [68]. <https://datacommons.org>
- [69]. India's wheat, rice production continue to reach record highs  
<https://www.world-grain.com/articles/13532-indias-wheat-rice-production-continue-to-reach-record-highs>
- [70]. Republic of India: Accelerating Agricultural Productivity Growth – World Bank Group, Agriculture 21 May 2014  
<https://openknowledge.worldbank.org/bitstream/handle/10986/18736/880930REVISED00ivity0Growth00PUBLIC.pdf?sequence=5&isAllowed=y>
- [71]. The future of food and agriculture Trends and challenges - Food and Agriculture Organization of the United Nations Rome, 2017  
<http://www.fao.org/3/i6583e/i6583e.pdf>
- [72]. Farmers' participation in agricultural development: The case of Fars province, Iran - Farshid Aref, Dept. of Soil Science, Science and Research Branch, Islamic Azad University, Fars, Iran. Indian Journal of Science and Technology, Vol. 4 No. 2 (Feb 2011) ISSN: 0974-6846  
[https://www.researchgate.net/publication/281641236\\_Farmers'\\_Participation\\_in\\_Agricultural\\_Development\\_The\\_Case\\_of\\_Fars\\_Province\\_Iran](https://www.researchgate.net/publication/281641236_Farmers'_Participation_in_Agricultural_Development_The_Case_of_Fars_Province_Iran)
- [73]. Integrated Weed Management Practices for Controlling Unwanted Vegetation By Lynn M. Sosnoskie -October 14, 2019; Organic Farmer  
<https://organicfarmermag.com/2019/10/integrated-weed-management-practices-for-controlling-unwanted-vegetation/>
- [74]. Studies of Morphological Characteristics and Production of Seeds Weeds of Species of Family Brassicaceae (Cruciferous) in Setifian High Plateau, Algeria - by Meriem Hani, Rafika Lebazda and Mohamed Fenni - Annual Research & Review in Biology 12(5): 1-9, 2017; Article no. ARRB.33473 ISSN: 2347-565X, NLM ID: 101632869  
<https://core.ac.uk/download/pdf/144780035.pdf>
- [75]. Review on Allelopathy of Exotic Invasive Plants – by Wang Chengxu, Zhu Mingxing, Chen Xuhui, Qu Bo, The Second SREE Conference on Chemical Engineering - Engineering 18 (2011) 240–246  
<https://reader.elsevier.com/reader/sd/pii/S1877705811028402?token=F0457D1A35518A2EC161D6AA788F76DC98EF2BCAB5559F570A571E69C436CD7284051EC3BD11084D0702373BACE4F78C&originRegion=eu-west-1&originCreation=20210730055011>
- [76]. Perennial weeds and their management. In: Weed Science and Management. Eds. N. T. Yaduraju, A. R. Sharma and T. K. Das, August 2020 - Publisher: Indian Society of Weed Science, ICAR-DWR, Jabalpur and Indian Society of Agronomy, ICAR-IARI, New Delhi  
[https://www.researchgate.net/publication/343382533\\_Perennial\\_weeds\\_and\\_their\\_management\\_InWeed\\_Science\\_and\\_Management\\_Eds\\_N\\_T\\_Yaduraju\\_A\\_R\\_Sharma\\_and\\_T\\_K\\_Das/stats](https://www.researchgate.net/publication/343382533_Perennial_weeds_and_their_management_InWeed_Science_and_Management_Eds_N_T_Yaduraju_A_R_Sharma_and_T_K_Das/stats)
- [77]. Calculation of LD50 values from the method of Miller and Tainter, 1944 – by Muhammad Akram Randhawa, Department of Pharmacology, College of Medicine, University of Dammam, Dammam, Saudi Arabia  
<https://www.ayubmed.edu.pk/JAMC/PAST/21-3/Randhawa.pdf>
- [78]. Environmental Chemistry and Ecotoxicology of Hazardous Heavy Metals: Environmental Persistence, Toxicity, and Bioaccumulation – Review article by Hazrat Ali, Ezzat Khan, and Ikram Ilahi - Journal of Chemistry, Volume 2019, Article ID 6730305, 14 pages, <https://doi.org/10.1155/2019/6730305>
- [79]. Why Regulators Lost Track and Control of Pesticide Risks: Lessons From the Case of Glyphosate-Based Herbicides and Genetically Engineered-Crop Technology Charles M. Benbrook - Current Environmental Health Reports volume 5, pages 387–395 (2018)  
<https://link.springer.com/article/10.1007/s40572-018-0207-y>
- [80]. The impact of foliar application of boron nano-chelated fertilizer and boric acid on fruit yield, oil content, and quality attributes in olive (*Olea europaea* L.) Zohre Rohi Vishekaia, Ali Soleimania, Esmaeil Fallahib, Mahmood Ghasemnezhad, Akbar Hasanid,  
<https://doi.org/10.1016/j.scienta.2019.108689>
- [81]. Foliar Application of Fertilizers by Wade W. McCall; Hawaii Cooperative Extension Service, College of Tropical Agriculture and Human Resources - University of Hawaii, General Home Garden Series no. 24  
<https://www.ctahr.hawaii.edu/oc/freepubs/pdf/GHGS-24.pdf>
- [82]. Integrated Weed Management Practices for Adoption in the Tropics - by Wendy-Ann P. Isaac, Puran Bridgemohan and Wayne G. Ganpat, 12 June 2013, DOI: 10.5772/55950  
<https://www.intechopen.com/chapters/44996>
- [83]. The integration of chemical and biological control of the spotted alfalfa aphid: Field experiments on the effects of insecticides by Vernon M. Stern, Robert van den Bosch - Hilgardia 29(2):103-130.



- DOI:10.3733/hilg.v29n02p103. October 1959.  
<http://hilgardia.ucanr.edu/Abstract/?a=hilg.v29n02p103>
- [84]. Field weed population dynamics: a review of model approaches and applications – by N Holst, I A Rasmussen, L Bastiaans - Department of Integrated Pest Management, Danish Institute of Agricultural Sciences, Flakkebjerg, Slagelse, Denmark, and Crop and Weed Ecology Group, Wageningen University, Wageningen, the Netherlands February 2007 - Weed Research 47(1) DOI: 10.1111/j.1365-3180.2007.00534.x  
[https://www.researchgate.net/publication/40110686\\_Field\\_weed\\_population\\_dynamics\\_A\\_review\\_of\\_model\\_approaches\\_and\\_applications](https://www.researchgate.net/publication/40110686_Field_weed_population_dynamics_A_review_of_model_approaches_and_applications)
- [85]. Weeds and Climate Change: Supporting weed management adaptation - Technical Guide by John K. Scott, Bruce L. Webber, Helen Murphy, Noboru Ota, Darren J. Kriticos, and Barton Loechele  
[https://adaptnrm.csiro.au/wp-content/uploads/2014/08/Adapt-NRM\\_M2\\_WeedsTechGuide\\_5.1\\_LR.pdf](https://adaptnrm.csiro.au/wp-content/uploads/2014/08/Adapt-NRM_M2_WeedsTechGuide_5.1_LR.pdf)
- [86]. Comparison of synthetic and organic herbicides applied banded for weed control in carrots (*Daucus carota* L.) – by D. C. Main, K. R. Sanderson, S. A. E. Fillmore, and J. A. Ivany, Canadian Journal of Plant Science - 27 June 2013, <https://doi.org/10.4141/cjps2013-052>
- [87]. Effect of Continuous Application of Herbicide on Soil and Environment with Crop Protection Machinery in Southern Adamawa State – by Daniel Mada, Duniya N, Idris G. Adams. International Refereed Journal of Engineering and Science (IRJES) ISSN (Online) 2319-183X, (Print) 2319-1821 Volume 2, Issue 6 (June 2013), PP.04-09  
<http://www.irjes.com/Papers/vol2-issue6/Version-2/B260409.pdf>
- [88]. Pesticide usage pattern in tea ecosystem, their retrospects and alternative measures - G. Gurusubramanian, A. Rahman, M. Sarmah, Somnath Ray and S. Bora - November 2008 Journal of Environmental Biology 29(6):813-26, [https://www.researchgate.net/publication/24211901\\_Pesticide\\_usage\\_pattern\\_in\\_tea\\_ecosystem\\_their\\_retr ospects\\_and\\_alternative\\_measures](https://www.researchgate.net/publication/24211901_Pesticide_usage_pattern_in_tea_ecosystem_their_retr ospects_and_alternative_measures)
- [89]. Pesticides, environment, and food safety – by Fernando P. Carvalho; 09 June 2017  
<https://onlinelibrary.wiley.com/doi/full/10.1002/fes3.108>
- [90]. From Smart Farming towards Agriculture 5.0: A Review on Crop Data Management – by Verónica Saiz-Rubio and Francisco Rovira-Más - Agricultural Robotics Laboratory (ARL), Universitat Politècnica de València, Camino de Vera, s/n. 46022 Valencia, Spain; Agronomy 2020, 10, 207; <https://www.mdpi.com/2073-4395/10/2/207/pdf>